

UNIVERSITATEA AGRARĂ DE STAT DIN MOLDOVA

ISSN 2587-3202 (electronic)

ISSN 1857-0003 (print)

**AGRICULTURAL SCIENCE
ȘTIINȚA AGRICOLĂ**

**Nr. 1
2022**

Chișinău 2022

CUPRINS

Pantelimon Borozan, Simion Musteața, Valentina Spînu, Alexei Spînu, Mihail Statnic Cercetări privind crearea liniilor consangvinizate și a hibrizilor de porumb timpuriu _____	3
Alexandru Rusu, Ludmila Bulat, Olga Arhip Asimilarea azotului și productivitatea culturilor în diverse procedee de fertilizare cu paie _____	12
Vasile Plămădeală, Ludmila Bulat, Natalia Bîstrova Bilanțul elementelor nutritive la aplicarea deșeurilor locale organogene pe cernoziom levigat _____	20
Алексей Постолати, Марина Рудой Корреляционный анализ сопряженности уровня урожая с биометрическими показателями и элементами его структуры у озимой пшеницы _____	27
Ananie Peșteanu, Andrei Lozan Influența regulatorilor de creștere asupra obținerii producțiilor înalte în plantațiile de cireș din soiul kordia altoite pe portaltolul maxma _____	32
Nadejda Mihnea, Diana Climăuțan, Cristian Roșca Reacția formelor parentale și hibrizilor f2 de tomate la temperaturi stresante _____	42
Татьяна Игнатьевна Щербакова Скрининг штаммов грибов trichoderma pers. для защиты растений от albifimbria verrucaria, ранее известного как myrothecium verrucaria _____	50
Irina Mihailov Staflinidofauna (coleoptera, staphylinidae) rezervației naturale de stat „Plaiul fagului” _____	57
Dionisie Boaghie Unele aspecte privind studiul creșterilor în diametru la goruneturile de proveniență vegetativă (lăstari) din republica Moldova _____	64
Sorin Macoviciuc, Constantin Bogdan Nechita, Ionel Bogdan Cioroiu, Valeriu Cotea, Marius Niculaua Effect of added sulphur dioxide levels on the aroma characteristics of wines from panciu wine region _____	73
Kryštof Mareš, Denisa Marešová, Tatiana Alexiou-Ivanova, Yayan Satyakti, Adrian Furculița Municipal solid waste management in Bandung, Indonesia: is incineration the right way to treat your waste? _____	78

CONTENTS

Charles Amarachi Ogbu, Marek Jelínek, Tatiana Alexiou-Ivanova, Iulia Corman The more wine, the more gas? estimation of the bioenergy potential of winery wastewater in moldova: contributions to sustainable development _____	85
Eduard Banari Evaluarea performanțelor ecologice ale motorului cu aprindere prin comprimare alimentat cu diverse tipuri de combustibili _____	92
Victor Popescu, Natalia Țislinscaia, Sergiu Popa, Vitali Vișanu, Mihail Balan, Mihail Melenciuc, Cristian Malai Sistem pentru deshidratarea fructelor cu eficiență energetică înaltă _____	97
Victor Popescu, Mihail Balan, Igor Beșleagă, Constantin Nesterenco, Tatiana Balan, Alexandru Dichii Studiul privind instalația pentru uscarea semințelor în strat suspendat _____	103
Tatiana Cușmenco Influența fructelor asupra proprietăților organoleptice ale iaurtului din lapte de capră și vacă _____	108
Valentina Stici, Tatiana Costin Definirea metodologiei spectrofotometrice uv-vis de evaluare a concentrației malondialdehidei ca marker al calității în trasabilitatea a trei tipuri de carne _____	116
Rita Golban Evaluări microbiologice ale unor sortimente de brânzeturi _____	125
Tatiana Golubenko, Olena Razanova Optimizing the use of protein in the young cattle body _____	143
Veronica Prisăcaru Engaging stakeholders in order to increase the quality of the agricultural higher education – current state and directions for improvement _____	153

Redactor-șef: Liviu VOLCONOVICI, doctor habilitat, profesor universitar (Republica Moldova)

Redactor-șef adjunct: Ilie CERCEL, doctor, conferențiar universitar (Republica Moldova)

Colegiul de redacție:

Iurie MELNIC, doctor, conferențiar universitar (Republica Moldova)

Vladimir BAUTIN, doctor habilitat, profesor universitar, academician al AȘR (Rusia)

Tlektes ESPOLOV, doctor habilitat, profesor universitar, academician al AȘK (Kazahstan)

Serghei KORLIUK, doctor habilitat, profesor universitar (Ucraina)

Arshaluys Poghos TARVERDYAN, doctor habilitat, profesor universitar, academician al AȘA (Armenia)

Vasile VÎNTU, doctor, profesor universitar (România)

Gerard JITĂREANU, doctor, profesor universitar (România)

Radu SESTRĂȘ, doctor, profesor universitar (România)

Florin STĂNICĂ, doctor, profesor universitar (România)

Victor ȘEREMET, doctor habilitat, profesor universitar (Republica Moldova)

Marian GRIGORE, doctor habilitat, profesor universitar (Republica Moldova)

Alexandru FRECĂUȚEANU, doctor habilitat, profesor universitar (Republica Moldova)

Gheorghe NICOLAESCU, doctor, conferențiar universitar (Republica Moldova)

Valerian BALAN, doctor habilitat, profesor universitar (Republica Moldova)

Ion BACEAN, doctor, conferențiar universitar (Republica Moldova)

Vladimir GOROBET, doctor, conferențiar universitar (Republica Moldova)

Oleg HORJAN, doctor, conferențiar universitar (Republica Moldova)

Mihail POPOVICI, doctor, conferențiar universitar (Republica Moldova)

Redactor-coordonator: Viorica LUPU

Revizuire text în limba engleză: Elena HODENCO

Revizuire text în limba rusă: Ana BRĂIESCU

DOI: <https://doi.org/10.55505/sa.2022.1.20>

UDC: 636.237.21.085.6

OPTIMIZING THE USE OF PROTEIN IN THE YOUNG CATTLE BODY

Tatiana GOLUBENKO, Olena RAZANOVA

Abstract. The organization of complete feed rations for cattle meeting the needs of animals in energy, protein, amino acids, minerals and vitamins in terms of basic nutrients and biologically active substances represents the main condition to achieve the genetic potential of productivity. The purpose of the paper was to optimize the protein utilization in the body of young cattle by synchronizing the processes of fermentation of nitrogen-containing substances and carbohydrates in feed. Barothermal treatment of concentrated feed mixtures with high protein degradability and a high content of non-structural carbohydrates causes a more uniform fermentation of nutrients, intensification of microbial protein synthesis processes and an increase in the efficiency of energy use of carbohydrates and protein nitrogen by rumen microflora. This is evidenced by an increase in the number of ciliates by 4.4–8.1% and a decrease in the concentration of ammonia and volatile fatty acids in the rumen fluid by 5.2–9.2 % and 3.4–4.3 %, respectively. A decrease in nitrogen losses is indicated by a decrease in the content of urea in animals' blood by 2.0–9.0%. Extrusion of concentrated feeds with high protein degradability and high content of non-structural carbohydrates improves animal productivity and feed efficiency. The average daily gain in live weight increases by 4.8–6.0% in the experimental group. As a result, feed consumption is reduced by 2.7–6.9%, while protein consumption is reduced by 2.6–5.7%.

Key words: Bulls; Feed; Extrusion; Barothermal treatment; Protein; Carbohydrates; Productivity.

INTRODUCTION

The organization of complete feed rations meeting the needs of animals in energy, protein, amino acids, minerals and vitamins in terms of basic nutrients and biologically active substances represents the main condition for achieving the genetic productivity potential (Skoromna, O.I. et al., 2019; Golubenko, T.L. 2018; Razanova, O.P. et al., 2022). There are certain feeding rules for each farm animal. Nutrition can be influence in different ways, i.e., to enhance or inhibit the conversion of certain substances during the metabolic activity, as well as to enhance their synthesis or decomposition by changing the amount and properties of feed and feeding conditions ensuring the formation of animal products (Skoromna, O.I. et al. 2018; Razanova, O.P. 2018; Grymak, Y. et al. 2020).

Protein is the most valuable component of feed; its level and quality determine animals' productivity. A complete protein nutrition of ruminants means meeting the needs of the animals' body for amino acids available for metabolism. However, the deficiency of feed protein and its irrational use makes the protein to be one of the most important limiting factors in the systems of intensive milk and meat production (Magowan, E. et al. 2010; De Oliveira, P.T. et al. 2011; Acedo, T.S. et al. 2011).

It is difficult and not cost-effective to obtain high productivity from animals by increasing the amount of high-protein feed in rations. It causes overconsumption of feed and an increase of the outputs cost. It also negatively affects the health of animals entailing a reduction of their productive use (Jeroch, H. 2008; Ahmad, M., Ali, A.A. 1994).

The modern concept of nutritional physiology is based on the fact that animal's need for protein is satisfied by the amino acids of the microbial protein and the protein that is not degraded in the rumen (Johansson, B. et al. 2012; Gurin, V.K. et al. 2016).

Creating conditions in the rumen that ensure maximum synthesis of microbial protein with a simultaneous increase in the flow of fodder protein into the intestine is one of the main conditions for the effective use of protein in the body. Microbial protein is unable to meet the increasing needs of the body for amino acids with an increase in the productivity of animals. In this case, the role of the transit fodder protein increases. Although, the higher the productivity of animals, the greater the contribution of the dietary protein not degraded in the rumen to the total pool of amino acids in the body. Thus, a high-quality protein for ruminants is a protein with a low degradability in the rumen, with a valuable amino acid composition and well digested in the intestine (Montiel, M.D. et al. 2012).

Protein degradation in the rumen is a complex, multiphase, enzymatic process influenced by many factors. Some of them are associated with the nature and properties of the protein i.e., its susceptibility

to fermentation, other factors are associated with the features of cicatricial digestion (pH, of microorganism species composition, etc.) (Bessarab, G.V. et al. 2012).

The degradation level and rate are important factors determining the efficiency of the microbial protein biosynthesis in the rumen, the general digestibility of nutrients, and the feed nitrogen utilization by animals (Koksal, B.H. et al. 2011; Wang, C. et al. 2009).

The efficiency of nitrogen utilization is highly dependent on the concentration of energy available for exchange, it implies significant fluctuations in the degradability of the crude protein of individual feeds. It seems relevant to research the degradability dynamics of the crude protein derived from forage products changing the specific energy content.

The proteins present in the rumen for a long period of time degrade more intensively than the proteins quickly evacuated from it. The protein of roughage is hidden under the cell membrane, rich in cellulose and lignin. That's why longer exposure to proteolytic enzymes is required for its degradability. Cereal feeds are faster taken from the proventriculus but they are degraded to a large extent indicating the physicochemical properties of their protein (Antonovich, A.M., Bessarab, G.V. 2018).

The ability to regulate the degree of protein degradability in the proventriculus represents an important issue in the nutrition of ruminants. As a rule, it is required to reduce the degradability of feed protein without drastic changes in its digestibility in the intestine. It can be achieved in two ways. The first one is the selection of natural feeds; their protein should be resistant to degradation in the rumen. This method has found relatively wide application, but it is not always possible because the range of feeds for ruminants is limited or not economically justified. It is possible to reduce the breakdown of dietary protein by physically or chemically affecting the protein in the feed. Considering the physical methods, we should mention that the most famous technique is exposure to high temperature. Such techniques contribute to the preservation of nutrients in feed, they also reduce the solubility and degradability of protein in them. Heat treatment of high-protein feed can reduce the solubility and disintegration of protein by 1.5 - 2 times (Bessarab, G.V. et al. 2018).

The purpose of the paper was to optimize the protein utilization in the body of young cattle by synchronizing the processes of fermentation of nitrogen-containing substances and carbohydrates in feed.

MATERIALS AND METHODS

The researches were carried out on the experimental animals: black-and-white bulls at the age of 3-6, 6-9, 9-12 and 12-18 months (Table 1).

The animals from the control group were fed the ground mix of concentrates, while the experimental group was fed the extruded mix of concentrates. The samples of rumen fluid were taken during the experiments and the degradability of the concentrated feed protein was researched. The hematological parameters and growth energy of experimental animals were determined.

Table 1. Feeding patters

Group	Number of animals, heads	Age of animals, months	Duration, days	Feeding characteristics
Experiment No. 1				
I control	3	3-6	60	BD (ground mix of concentrates)
II experimental	3	3-6	60	BD (extruded mix of concentrates)
Experiment No. 2				
I control	3	6-9	60	BD (ground mix of concentrates)
II experimental	3	6-9	60	BD (extruded mix of concentrates)
Experiment No. 3				
I control	3	9-12	60	BD (ground mix of concentrates)
II experimental	3	9-12	60	BD (extruded mix of concentrates)
Experiment No. 4				
I control	3	12-18	60	BD (ground mix of concentrates)
II experimental	3	12-18	60	BD (extruded mix of concentrates)

Chemical composition of the feed used in the experiments was determined according to the scheme

of general zootechnical analysis. Therefore, the initial, hygroscopic and total moisture, mass fraction of crude protein were determined using the automatic analyzer UDK132 and UDK159 (VELP, Italy), the mass fraction of crude fiber using a semi-automatic FIWE-6 analyzer, the mass fraction of crude fat, mass fraction of crude ash and organic matter were determined in the feed according to the generally accepted zootechnical methods (Malchevskaya, E.N., Milenkaya, G.S. 1981; Petukhova, E.A. et al. 1989).

The intensity of bulls' cicatricial digestion was studied by sampling the liquid part of the rumen contents through the fistula 2-2.5 hours after morning feeding and it was filtered through four layers of gauze.

The following parameters were determined e liquid part of the of the cicatricial content, i.e., the concentration of hydrogen ions (pH), ammonia concentration and total nitrogen using the automatic analyzer UDK132 and UDK159 (VELP, Italy), the total amount of VFA by steam distillation in Markgam apparatus, the number of ciliates by counting in a 4-mesh Goryaev chamber (Kurilov, N.V. et al. 1987).

Feed protein degradability was determined according to GOST 28075-89. Samples of concentrated feed were placed in nylon bags. The incubation period of the researched concentrated feed in the rumen was 2, 4, 6, 8 and 12 hours.

We have also researched the feed intake by carrying out ten-day control feedings for two adjacent days according to the difference in weight of the given feed and uneaten residues, the growth rate and average daily gain in live weight of animals by individual weighing at the beginning and at the end of the experiment, as well as the efficiency of feed utilization by calculating energy and protein consumption.

Statistical evaluation of differences was performed using Student's t-test [30]. We proceeded depending on the volume of the analyzed material when evaluating the values of the reliability criterion. The differences were considered significant at $P < 0.05$.

RESULTS AND DISCUSSIONS

The experimental animals received a diet consisting of corn silage and compound feed in the first and second experiments. The control group was additionally fed by a mixture of ground barley grains and field peas, and the experimental group was additionally fed by an extruded mixture (Table 2).

The share of concentrated feed accounted for 36% in the diet structure. Herbal fodder accounted for 64% in the diet structure. The consumption of corn silage increased by 2.2% in the experimental group. The animals ate the concentrated feed completely.

The experimental young animals consumed 4.3-4.4 kg per head of dry matter per day. The metabolic energy content was 10.1 MJ/kg in 1 kg of dry matter of the diet in the experimental group. The share of crude protein accounted for 11.9%. Protein degradability of the control group was 80%, in the experimental one it was 76%. The amount of fiber in dry matter was 26%.

The effect of feeding an extruded mixture of concentrates with a high content of digestible protein and non-structural carbohydrates on indicators of cicatricial digestion was researched in the second physiological experiment. As a result, the regularity of the digestion processes in the rumen of 6-9-month-old young cattle was investigated. The share of concentrated feed was 38.4-39.5% in the diet structure. Herbal fodder accounted for 60.5-61.6%. The animals ate the concentrated feed completely. An increase of 4.9% in the consumption of corn silage was recorded in the experimental group. The experimental young animals received 6.21-6.43 kg/head of dry matter daily. The content of metabolizable energy in the dry matter of the diet of the experimental groups was 9.9 MJ/kg. The share of crude protein in dry matter was 12.6-12.8%, fiber share was 27%. According to research data, protein degradability in the corn silage was 75.6%, in the compound feed - 81%, in the mixture of ground barley and field peas - 82.1% and in the extruded mixture of barley and field peas it was of 53.8%. Thus, extrusion helped to reduce the disintegration of the grain mixture by 28.3%.

Total nitrogen content was by 8.3% higher, and the ammonia content was by 8.7% lower in the rumen of the 3-6-month-old animals fed extruded grain mixture (Table 3).

Table 2. Diet of 3-9-month-old experimental animals

Feed and nutrients	Age, months			
	3-6		6-9	
	Group			
	I	II	I	II
Corn silage, kg	9.2	9.4	14.20	14.90
Compound feed KR-3, kg	1.0	1.0	1.4	1.4
Ground grain mixture, kg	0.5		0.6	
Extruded grain mixture, kg		0.5		0.6
Diet components				
Feed units	4.48	4.58	5.89	6.09
Exchange energy, MJ	43.7	44.5	61.4	63.6
Dry matter, kg	4321	4397	6.21	6.43
Crude protein, g	518	527	795	816
Degradable protein, g	415	399	618	588
Non-degradable protein, g	104	128	177	228
Crude fat, g	225	230.0	230	240
Crude fiber, g	1,117	1,140	1,687	1,764
NFE, g	204	2,090	3,113	3,225
Calcium, g	28.8	29.3	41.0	42.6
Phosphorus, g	18.5	18.8	26.4	27.3
Magnesium, g	10.23	10.4	15.4	16.0
Potassium, g	55.13	56.2	84.1	87.6
Sulphur, g	8.59	8.74	12.9	13.5
Iron, mg	1,173	1,197	1,781	1,865
Copper, mg	88.5	88.9	98.4	99.6
Zinc, mg	189	191	266	275
Manganese, mg	329	334	468	486
Cobalt, mg	1.50	1.50	1.67	1.69
Iodine, mg	1.80	1.83	2.61	2.71

Table 3. Parameters of cicatricial digestion of 3-9-month-old animals

Indicator	Age, months			
	3-6		6-9	
	Group			
	I	II	I	II
pH	6.04±0.16	6.18±0.18	6.46±0.18	6.61±0.18
Volatile fatty acids, mmol/100 ml	10.6±0.40	10.23±0.18	10.73±0.28	10.36±0.21
Total nitrogen, mg/100 ml	134.5±14.5	145.7±14.89	100.3±1.95	103±1.05
Ammonia, mg/100 ml	13.8±0.6	12.6±0.40	13.35±0.65	12.27±0.74
Ciliates, thousand/ml	799±13.5	833±21.8	648±23.5	683±14.75

Ammonia level decrease and total protein level increase indicate an increase in the microbial protein synthesis due to the creation of more favorable conditions for the vital activity of microflora, its number increased by 4.4% in group II of 3-6-month-old animals. There was a decrease of 3.5% in the experimental group in the amount of volatile fatty acids. The reaction of the rumen pH did not change significantly reaching the level of 6.0-6.2 in all groups.

The pH level was slightly lower in the control group of 6-9-month-old animals being of 6.46. This indicator was 6.61 in the experimental group because of a decrease in the level of volatile fatty acids by 3.4%. There was a decrease in ammonia concentration of 8.1% in the experimental group. The increase in the total nitrogen amount and ammonia decrease may indicate an increase in the intensity of microbial protein synthesis due to a more uniform supply of nutrients to the rumen and the creation of favorable conditions for the vital activity of microflora, as evidenced by an increase in the number of ciliates by 5.4%.

However, all indicators were within the normal range despite some changes during the digestion processes in the rumen of animals.

According to research data, the hematological parameters were within the physiological norms (Table 4).

Table 4. Hematological parameters of 3-9-month-old calves

Indicator	Age, months			
	3-6		6-9	
	Group			
	I	II	I	II
Erythrocytes, 10 ¹² /l	6.24±0.13	6.49±0.12	6.74±0.13	6.99±0.120
Leukocytes, 10 ⁹ /l	10.05±0.25	10.23±0.49	10.55±0.25	10.33±0.230
Hemoglobin, g/l	106.1±6.3	110.2±4.51	108.55±5.95	109.17±6.060
Total protein, g/l	75.75±2.25	78.77±1.56	77.8±2.3	81.87±1.620
Glucose, mmol/l	2.49±0.16	2.33±0.03	2.82±0.26	2.81±0.050
Urea, mmol/l	4.1±0.14	4.02±0.14	4.72±0.16	4.33±0.150
Calcium, mmol/l	2.82±0.12	2.64±0.06	2.93±0.125	2.74±0.0670
Phosphorus, mmol/l	1.59±0.15	1.66±0.05	1.54±0.1	1.66±0.050
Hematocrit, %	34.55±1.85	34.73±1.22	32.4±0.8	33.83±0.6740

The 3-6-month-old experimental animals recorded an increase in the number of erythrocytes by 4.0 %, hemoglobin by 3.9 %, total protein by 4.0 % and phosphorus by 4.4 %. The level of glucose decreased by 6.4%, urea decreased by 2.0 % and calcium decreased by 6.4%. However, the noted differences were not significant.

The extruded mixture application influenced the blood composition of 6-9-month-old animals. Thus, young experimental animals had an increase in the content of erythrocytes by 3.7%, total protein by 5.2 %, phosphorus by 7.8%, hematocrit by 4.4%, urea by 8.3%, and calcium by 6.5%. However, the differences noted are not significant.

The control weighing of animals was carried out at the beginning and at the end of the experiment in order to control the live weight. Also, it was investigated the effect of extruded grain mixture on the productivity of animals (Table 5).

Table 5. Dynamics of live weight and efficiency of feed utilization by 3-9-month-old experimental young animals

Indicator	Age, months			
	3-6		6-9	
	Group			
	I	II	I	II
Live weight, kg:				
at the beginning of the experiment	132.7±1.3	133.1±1.80	205.2±1.6	204.5±1.70
at the end of the experiment	178.3±3.5	181.3±2.40	252.2±0.8	254.3±1.80
Gross growth, kg	45.6±2.2	48.2±10	47±2.4	49.8±2.80
Average daily gain, g	760±37	803.3±17.7	783±40	830±46.80
% to control	100	105.7	100	106.0
Feed consumption per 1 kg gain, feed units	5.89	5.70	7.52	7.34
% to control	100	96.8	100	97.3
Protein consumption per 1 kg of gain, kg	0.68	0.66	0.98	0.95
% to control	100	97.1	100	97.4

According to weighing data, feeding an extruded mixture of field peas and barley instead of ground grain caused an increase in the average daily gain of 3-6-month-old animals. Higher gains were recorded by group II, 804 g per day, which is by 5.8% higher than in group I. Feed consumption in this group decreased by 3.2% being of 5.7 feed units. The feed protein efficiency also increased by 2.9%.

According to our research data, higher gain was registered in the experimental group formed by 6-9-month-old animals. It was of 830 g per day or by 6.0% higher than in the control group. In this

group, the feed consumption was by 2.7% lower than in group I, the efficiency of feed protein utilization has also increased by 2.6%.

The dependence of the cicatricial digestion indicators of 9-12-month-old cattle was researched in the third experiment, when the animals were fed an extruded mixture of high-protein and high-carbohydrate concentrates. The effectiveness of the nutrient utilization in the body of animals was researched.

The diet of 9-18-month-old experimental animals consisted of corn silage, compound feed and a mixture of cereals and legumes. Also, the animals received plenty of silage. The control animals received a mixture of ground barley and field peas, and the experimental group animals received an extruded one (Table 6).

Table 6. Diet of 9-18-month-old experimental animals

Feed and nutrients	Age, months			
	9-12		12-18	
	Group			
	I	II	I	II
Corn silage, kg	16.20	16.50	17.30	17.60
Compound feed KR-3, kg	1.4	1.4	1.6	1.6
Ground grain mixture, kg	0.6		0.7	
Extruded grain mixture, kg		0.6		0.7
Diet components				
Feed units	7.36	7.48	8.46	8.58
Exchange energy, MJ	73.6	74.7	86.6	87.9
Dry matter, kg	7.16	7.27	9.41	9.55
Crude protein, g	862	863	958	976
Degradable protein, g	656	612	679	662
Non-degradable protein, g	206	251	279	314
Crude fat, g	254	260	220	223
Crude fiber, g	1619	1647	2533	2576
NFE, g	4002	4062	5166.0	5237.8
Calcium, g	45.56	46.28	62.3	63.2
Phosphorus, g	28.79	29.20	25.4	25.7
Magnesium, g	17.18	17.47	18.6	18.8
Potassium, g	94.05	95.67	173	176
Sulfur, g	14.34	14.58	17.1	17.4
Iron, mg	2021	2057	3952	4019
Copper, mg	102	102	151	151
Zinc, mg	291	295	291	295
Manganese, mg	521	529	626	635
Cobalt, mg	1.73	1.74	2.99	3.00
Iodine, mg	2.89	2.93	2.52	2.55

The share of concentrated feed for 9-12-month-old animals was 31%, the share of herbal feed was 69%. The animals ate the concentrated feed completely. There was a slight increase in the consumption of corn silage by the experimental group animals. The experimental young animals received daily 7.16-7.27 kg per head of dry matter. The content of metabolizable energy in the dry matter was 10.3 MJ per kg. The share of crude protein in dry matter was 11.4-11.6% and fiber share was 22.6%. Researchers have found that protein degradability was 76% in the control group and 71% in the experimental one. A decrease in the degree of protein degradation occurred because of the extruded concentrates feeding. Thus, the degradability of the protein derived from ground barley and field peas mixture is 82%, while the degradability of the protein derived from extruded barley and field peas mixture is 54%.

Then, the dependence of the indicators of protein metabolism and cicatricial digestion was researched when concentrated feeds were included in the diet, with a high degree of protein and carbohydrate breakdown, subjected to barothermal treatment, carried out on 12-18-month-old calves. The share of

concentrated feed was 33% and the share of herbal feed was 67%. The animals consumed the concentrated feed completely. There was a slight increase of haylage consumption by 1.7% in group II. The experimental young animals received 9.38-9.53 kg per head of dry matter, consequently 1 kg contained 9.23 MJ of metabolic energy, 102 g of crude protein and 270 g of fiber. The degradability of the ground grain mixture protein was 84%, the degradability of the extruded grain mixture protein was 58%. It decreased by 3.1% because of replacing the ground pea and barley mixture with extruded one.

According to research data, there was a decrease of the volatile fatty acids level by 4.3% and ammonia concentration decrease by 5.2% in the rumen fluid of 9-12-month-old calves from the experimental group (Table 7).

The decrease of an ammonia level and increase of total nitrogen indicate an intensification of the microbial protein synthesis due to a more uniform supply of nutrients to the rumen and the creation of more favorable conditions for the vital activity of microflora, which is indicated by an increase in the number of ciliates by 8.1% ($P < 0, 05$).

Table 7. Parameters of cicatricial digestion of 9-18-month-old animals

Indicator	Age, months			
	9-12		12-18	
	Group			
	I	II	I	II
pH	6.67±0.16	6.75±0.080	6.82±0.15	6.86±0.17
Volatile fatty acids, mmol / 100 ml	10.17±0.95	9.73±0.390	10.1±0.15	9.7±0.24
Total nitrogen, mg / 100 ml	97.6±5.65	97±0.9020	118.6±4.20	127.7±3.38
Ammonia, mg / 100 ml	14±0.4	13.27±0.2960	13.03±0.26	11.83±0.49
Ciliates, thousand / ml	750.5±7.5	811.33±9.61*	810.33±9.5	874±17.7*

According to research data on 12-18-month-old animals, the use of extruded mixture caused a volatile fatty acids level decrease by 4%, ammonia level decrease by 9.2%, total nitrogen increases by 7.7 % and ciliates increase by 7.9% ($P < 0.05$) in the rumen (Table 6).

Ammonia level decrease and total protein increase indicate that the intensity of microbial protein synthesis has enhanced because of a more uniform release of protein nitrogen and energy from the extruded mixture into the rumen fluid and the creation of more favorable conditions for the vital activity of microflora. Despite some changes in the digestion processes in the animals' rumen, all indicators were within the normal range.

Extruded mixture feeding had an effect on the blood composition of animals, but the indicators were within the physiological norms (Table 8).

Table 8. Hematological parameters of 12-18-month-old calves

Indicator	Age, months			
	9-12		12-18	
	Group			
	I	II	I	II
Erythrocytes, 1012/l	6.32±0.31	6.36±0.20	6.82±0.15	6.86±0.17
Leukocytes, 109/l	10.84±0.38	10.74±0.23	6.41±0.35	6.4±0.20
Hemoglobin, g / l	117.9±2.75	118.5±1.74	111.7±3.71	114.3±2.03
Total protein, g / l	73.5±2.3	74.23±1.64	73.8±1.56	77.4±1.81
Glucose, mmol / l	2.52±0.06	2.53±0.10	2.56±0.13	2.63±0.11
Urea, mmol / l	4.28±0.1	3.98±0.13	4.09±0.10	3.72±0.21
Calcium, mmol / l	2.6±0.13	2.66±0.10	2.66±0.06	2.75±0.02
Phosphorus, mmol / l	1.75±0.07	1.69±0.07	1.61±0.07	1.58±0.04
Hematocrit, %	32.67±2.515	33.55±0.97	34.33±1.42	35.13±1.14

There was a hematocrit increase by 2.7%, a urea decreases by 7.0 %, and a phosphorus decrease by 3.4% in the blood of 9-12-month-old animals of the experimental group. However, the noted differences are not significant.

There was an increase in the amount of total protein by 4.8%, glucose by 2.7 %, calcium by 3.4 %, hematocrit by 2.3% and a decrease in urea by 9%, phosphorus by 1.9% in the experimental group, and namely in the group of 12-18-month-old animals. However, the noted differences are not significant.

Feeding an extruded mixture of field pea and barley grains instead of ground grain caused an increase in growth energy and the efficiency of feed utilization (Table 9).

There was an increase in the average daily gain in live weight by 4.8% and a decrease in feed consumption by 6.9% in the 9-12-month-old animals from the experimental group. The feed protein efficiency also increased by 5.7%.

Table 9. Dynamics of live weight and efficiency of feed utilization by 12-18-month-old experimental young animals

Indicator	Age, months			
	3-6		6-9	
	Group			
	I	II	I	II
Live weight, kg:				
at the beginning of the experiment	132.7±1.3	133.1±1.80	205.2±1.6	204.5±1.70
at the end of the experiment	178.3±3.5	181.3±2.40	252.2±0.8	254.3±1.80
Gross growth, kg	45.6±2.2	48.2±10	47±2.4	49.8±2.80
Average daily gain, g	760±37	803.3±17.7	783±40	830±46.80
% to control	100	105.7	100	106.0
Feed consumption per 1 kg gain, feed units	5.89	5.70	7.52	7.34
% to control	100	96.8	100	97.3
Protein consumption per 1 kg of gain, kg	0.68	0.66	0.98	0.95
% to control	100	97.1	100	97.4

According to the results of weighing the 12-18-month-old animals, the higher gains were noted in the experimental group, and namely of 865 g per day, which is by 5.1% higher than in the control group. In the experimental group the feed consumption was lower by 3.6% than in the control one, the efficiency of feed protein utilization increased by 3.5%.

CONCLUSIONS

Extrusion of concentrated feed reduces the degradability of protein concentrates. The number of ciliates in the rumen fluid increases by 5.4-8.1%, total nitrogen increases by 8.3%, and the concentration of ammonia and volatile fatty acids decreases by 6.2-9.2% and 3.4-4.3%, respectively. Thus, the processing of concentrates stimulates the development of proventriculus microflora and reduces protein loss in the rumen.

Barothermal treatment of concentrates with a high protein degradability and a high content of non-structural carbohydrates increases the productivity of animals and the efficiency of feed utilization. The average daily gain in live weight of animals from the experimental group increased by 4.8-6.0%. As a result, feed consumption is reduced by 2.7-6.9%, protein consumption is reduced by 2.6-5.7%.

REFERENCES

1. ACEDO, T.S. et al. (2011). Protein sources in supplements for bulls in the dry-rainy transition season: nutritional characteristics. In: Arquivo Brasileiro de Medicina Veterinária e Zootecnia, № 63(4), pp. 895-904. Available: DOI:10.1590/S0102-09352011000400015
2. AHMAD, M., ALI, A.A. (1994). Growth rate in buffalo and Sahiwal calves under optimum feeding conditions. In: 15 Annual Rep., LPRI, Bahadar Nagar, Okara, Pakistan, pp. 9-60.
3. ANTONOVICH, A.M., & BESSARAB, G.V. (2018). Cicatricial digestion and protein breakdown of high protein feeds in the rumen, depending on the processing method. In: Modern technologies of agricultural production, pp. 118-120.

4. BESSARAB G.V. et al. (2018). The effectiveness of different methods of preparing grain for feeding. In: Current nutritional technology products, pp. 123-127.
5. BESSARAB, G.V. et al. (2019). Increasing the efficiency of raising young cattle by rationing the protein in the diet (Increasing the efficiency of growing young cattle by rationing the digestible protein in the diet). In: Modernization of agricultural education: integration of science and practice. Collection of scientific papers based on the materials of the V International Scientific and Practical Conference, pp. 212-215.
6. De OLIVEIRA, P.T. et al. (2011). Physiological responses and performance of sheep on pasture supplemented with different protein sources. In: Rev. Ceres, № 58(2), pp. 185-192.
7. GOLUBENKO, T.L. (2018). Nutritional value of veal as the most important criterion for evaluating raw meat for baby food. In: Ukrainian Journal of Ecology, № 8(1), pp. 637-643. Available: DOI:10.15421/2018_260
8. GRYMAK, Y. et al. (2020). Influence of “Thireomagnile” and “Thyrioton” preparations on the anti-oxidant status of pregnant cows with symptoms of endotoxiosis. In: Ukrainian Journal of Ecology, № 10(1), pp. 122-126. Available: DOI:10.15421/2020_19
9. GURIN, V.K. et al. (2016). Conversion of feed by breeding gobies into products when feeding diets with different protein quality. In: Zootechnical science of Belarus, № 51(1), pp. 257-266.
10. JEROCH, H. (2008). The significance of rapeseed and rapeseed products for animal nutrition and the quality of animal products. In: Zemes ukio mokslai, № 15(4), pp. 40-52.
11. JOHANSSON, B. et al. (2012). Cold-pressed rapeseed cake or rapeseed to dairy cows - milk production and profitability. In: Tacking the Future Challenges of Organic Animal Husbandry: Organic Animal Husbandry Conference, Hamburg, 12-14 Sept. 2012. Available: DOI:10.1007/s13165-014-0094-y.
12. KOKSAL, B.H. et al. (2011). Rumen degradation characteristics of glucose-treated canola meal and canola seed. In: Veterinarija ir zootechnika, № 56(78), pp. 73-79.
13. KURILOV, N.V., SEVASTYANOVA, H.A., KORSHUNOV, V.N. et al. (1987). The Study of Digestion in Ruminants: Method. VASHNIL, Research Institute of Physiology, Biochemistry and nutrition of agricultural animals. Bobrovsky: VNIIFBIP agricultural animals, p. 104.
14. MAGOWAN, E. et al. (2010). The effect of supplementary grass silage and standard concentrate on milk fat fatty acid composition and iodine value when cows are fed a whole rapeseed-based concentrate at pasture. In: Irish Journal of Agricultural and Food Research, № 49(2), pp. 129-139. Available: DOI:10.2307/41219178.
15. MALCHEVSKAYA, E.N., Milenkaya, G.S. (1981). Quality assessment and analysis of zootechnical feed. Minsk: Uradzhaj.
16. MONTIEL, M.D. et al. (2012). Deactivation of tannin in high-moisture sorghum grain with polyethylene glycol or urea. In: Archivos de Zootecnia, № 61(234), pp. 235-244. Available: DOI:10.4321/S0004-05922012000200008
17. PETUKHOVA, E. A. et al. (1989). Zootechnical analysis of feed. Moscow: Agropromizdat. [in Russian].
18. RAZANOVA, O.P. (2018). Increasing meat quality quails fed by biological active additives based on submerged bees. In: Ukrainian Journal of Ecology, № 8(1), pp. 631-636. Available: DOI:10.15421/2017_259
19. RAZANOVA, O.P. et al. (2022). Influence of PVMS Intermix on living weight and linear measurements of bulls of Ukrainian black and spinned milk breed. Bulletin of Sumy National Agrarian University. The series: Livestock, № 1(48), pp. 65-71. DOI: 10.32845/bsnau.lvst.2022.1.10
20. ROKITSKY, P.F. (1973). Biological statistics. Minsk, 320 p. [in Russian]
21. SKOROMNA, O.I. & RAZANOVA, O.P. & TKACHENKO, T.Y. (2019). Effect of lysine feeding allowance on growth performance and carcass characteristics of growing pigs. In: Ukrainian Journal of Ecology, № 9(4), pp. 646-650. Available: DOI:10.15421/2019_803
22. SKOROMNA, O.I. et al. (2018). Balancing ration of dairy cows on calcium, phosphorus and iron indices for milk production and exchange processes in the organism. In: Ukrainian Journal of Ecology, № 8(3), pp. 92-97.

23. WANG, C. et al. (2009). Effects of citric acid supplementation on rumen fermentation, urinary excretion of purine derivatives and feed digestibility in steers. In: Journal of the Food and Agriculture, № 89(13), pp. 2302-2307. Available: <https://doi.org/10.1002/jsfa.3724>

INFORMATION ABOUT AUTHORS

GOLUBENKO Tatiana  <https://orcid.org/0000-0003-1427-3950>

PhD in Agricultural Sciences, Faculty of Technology of Production and Processing of Livestock and Veterinary Products, Vinnytsia National Agrarian University, Ukraine

E-mail: aponas504@ukr.net

RAZANOVA Olena  <https://orcid.org/0000-0001-5552-9356>

PhD in Agricultural Sciences, Faculty of Technology of Production and Processing of Livestock and Veterinary Products, Vinnytsia National Agrarian University, Ukraine

E-mail: olenaop0205@ukr.net

Data prezentării articolului: 01.04.2022

Data acceptării articolului: 04.05.2022