

# International Journal of Agriculture and Food Science



International Journal of Agriculture and Food Science is a peer reviewed, indexed, refereed and open access journal. Considers review and research articles related to agricultural economics, agro-ecosystems, Crop physiology, crop production and management, agro-climatology and agronomic modeling, farmers psychology, plant-soil relationships, crop quality and post-harvest physiology, Cropping of sustainable development, farming and cropping systems, plant protection and nutrition, genetic and breeding resources, storage technologies, etc.

Editor-in-Chief: Dr. Manas Mohan Adhikary

## Editorial Board

ISSN: P-ISSN: 2664-844X, E-ISSN: 2664-8458

Abbreviated Title: Int. J. Agric. Food Sci.

Impact Factor: 5.6

DOI: <https://dx.doi.org/10.33545/2664844X>

The journal provides the choice of both open access and subscription mode of publication to the authors and publishes almost all types of write-ups like research articles, reviews, case reports, case studies, commentary, letters to the editor, mini-reviews, opinions, short communication, book review, editorials, etc.

All the articles submitted to the journal are suitably reviewed by experts and the honorary editorial board. Prospective authors can submit their manuscripts at [agri.submit@gmail.com](mailto:agri.submit@gmail.com). Read Instructions to Author for more details.

## The Journal is Indexed and Abstracted in the Following Databases



## Editorial Board

### Editor in Chief



**Dr. Manas Mohan Adhikary**  
Professor  
Department of Agricultural Extension, Bidhan Chandra Krishi Viswavidyalaya, West Bengal, India  
Email: dradhikary@gmail.com

### Editors



**Dr. Dipan Adhikari (Ph.D.)**  
Assistant Professor  
Department of Botany, Hooghly Mohsin College Chinsurah, Hooghly, West Bengal, India  
Email: dipanadhikari@gmail.com



**Dr. J. John Peter Paul (M.Sc., M.Phil., Ph.D.)**  
Assistant Professor  
Department of Botany, St. Xavier & College (Autonomous), Palayamkottai, Tamil Nadu, India  
Email: johnarock2008@yahoo.com  
Phone: 9442955038



**Rahman Ahm Mahbubur**  
Associate Professor  
Department of Botany, University of Rajshahi, Rajshahi, Bangladesh



**Dr. Z N Tahmida Begum**  
Associate Professor  
Department of Botany, University of Dhaka, Dhaka, Bangladesh



**Dr. Nasrullah**  
Assistant Professor  
Department of Botany, University of Malakand, Pakistan



**Dr. Manzoor Hussain (PhD)**  
Professor  
Department of Botany, Hazara University, Mansehra, Khyber, Pakhtunkhwa, Pakistan



**Weili Yang**  
Professor  
Department of Plant Biology, Michigan State University, USA








**Chenping Xu**  
Assistant Professor  
Department of Plant Biology and Pathology, Rutgers University, USA



**Dr. Phil Harris**  
Assistant Professor  
Department of Plant Sciences, School of Science and The Environment, Coventry University, UK



**Catherine Hyland**  
Assistant Professor  
Department of Annals of Botany, Department of Plant Sciences, University of Oxford, South Parks Road Oxford, UK

	<p><b>Dr. Tarek Kapiel</b>  Assistant Professor  Plant Biotechnology at Biology Department, Faculty of Sciences, Al-Baha University, Saudi Arabia</p>
	<p><b>Dr. Tanweer Azam</b>  Professor  Ambo University, Ethiopia</p>
	<p><b>Siraj Mammo Yimer</b>  Professor  Ambo University, Ethiopia</p>
	<p><b>Chan Lai Kengm</b>  Professor  Department of Plant Tissue and Cell Culture Laboratory, University of Science, Malaysia</p>
	<p><b>Dr. Fahrul Zaman Huyop</b>  Professor  University of Technology, Malaysia</p>
	<p><b>Gurleen Sidhu</b>  Professor  Plant breeding and Genetics, University of Guelph, Guelph, Canada</p>
	<p><b>Gale G Bozzo</b>  Associate Professor  Department of Plant Agriculture, University of Guelph, Guelph, Canada</p>
	<p><b>Mohamed Mahgoub Azooz</b>  Professor  Department of Botany, Faculty of Science, South Valley University, Egypt</p>
	<p><b>Dr. Eman Abdel-Samiaa Ali Hussein Alam</b>  Professor  Department of Botany, Faculty of Science, Helwan University, Egypt</p>
	<p><b>Ngwengeh Brendaline Beloke</b>  Professor  University of Buea Cameroon, Cameroon</p>
	<p><b>Akwene Godwill Chenyuei</b>  Assistant Professor  University of Yaounde Cameroon, Cameroon</p>
	<p><b>Wenjuan Zhang (PhD)</b>  Assistant Professor  Xian Polytechnic University, China</p>
	<p><b>Dr. Alison Kim Shan Wee</b>  Assistant Professor  Guangxi University, China</p>
	<p><b>Dr. Nobuhiro Suzuki Website</b>  Professor  Sophia University, Tokyo, Japan</p>

	<p><b>Dr. Akira Kawabe</b>  Professor  Kyoto Sangyo University, Japan</p>
	<p><b>John Baling Fom</b>  Professor  University of Jos Nigeria, Nigeria</p>
	<p><b>Chinelo Ezeabara</b>  Professor  Department of Botany, Faculty of Biosciences, Nnamdi Azikiwe University, Awka, Anambra, Nigeria</p>
	<p><b>Dr. Kenneth Ekwealor (Ph.D)</b>  Senior Lecturer  Nnamdi Azikiwe University, Awka, Anambra, Nigeria  Email: drjckonkwo@yahoo.com  Phone: +2348035414358</p>
	<p><b>Saad Farouk Mohamed Hussien (PhD)</b>  Associate Professor  Agricultural Botany Department Faculty of Agriculture Mansoura university Mansoura Egypt  Email: gadalla@mans.edu.eg  Phone: 00201001105095</p>
	<p><b>Yao Saraka Didier Martial (Ph.D)</b>  Associate Professor &amp; Head  Ufr Sciences Biologiques, University of Peleforo Gon Coulibaly (UPGC), Korhogo, Côte D'Ivoire  Email: didierys@yahoo.fr  Phone: +22504737926</p>
	<p><b>Minhajur Rahman (MS in Botany)</b>  Assistant Professor  Department of Botany, University of Chittagong Chittagong, Bangladesh  Email: minhaj_bot@cu.ac.bd  Phone: +8801819624004</p>
	<p><b>Ahmed Jibrin Uttu (Ph.D)</b>  Associate Professor  Federal University Gashua, Nigeria  Email: jibuttu@yahoo.com  Phone: 08091066456</p>
	<p><b>Dr. Felix-Gastelum Ruben</b>  Professor  Autonomous University of Occident, Mexico</p>
	<p><b>José Lpez-Bucio</b>  Professor  Michoacan University of San Nicolás de Hidalgo, Mexico</p>
	<p><b>Chi-Yu Chen</b>  Associate Professor  National Chung-Hsing University, Taiwan</p>
	<p><b>Dr. Biing T Guan</b>  Associate Professor  National Taiwan University, Taiwan</p>

	<p><b>Bhisma Murti</b>  Professor  Sebelas Maret University, Indonesia</p>
	<p><b>Dr. Suardi Abang Abang</b>  Professor  University of Jakarta Indonesia, Indonesia</p>
	<p><b>Dr. Emmie M Cabanlit</b>  Associate Professor  University of Southeastern, Philippines</p>
	<p><b>Dr. Pablito Macapugay Magdalit</b>  Professor  College of Agriculture, University Philippines, Philippines</p>
	<p><b>Dr. Langa Tembo</b>  Professor  Department of Plant Science, University of Zambia, Lusaka, Zambia</p>
	<p><b>Christopher Phiri</b>  Professor  Department of Plant Science, University of Zambia, Lusaka, Zambia</p>
	<p><b>Rlc Wijesundera</b>  Associate Professor  Department of Plant Sciences, Faculty of Science, University of Colombo, Colombo, Sri Lanka</p>
	<p><b>Therese Lakshmi Shamala Tirimanne</b>  Associate Professor  Department of Plant Sciences, University of Colombo, Sri Lanka</p>
	<p><b>Dr. Nghia Thi Ai Nguyen</b>  Professor  Vietnam National University of Agriculture, Vietnam</p>
	<p><b>Van Tuan Tran</b>  Professor  Vietnam National University, Hanoi, Vietnam</p>
	<p><b>Dr. Ir Sim Kok Swee</b>  Professor  National University of Singapore, Singapore</p>
	<p><b>Dr. Gan-Moog Chow</b>  Associate Professor  National University of Singapore, Republic of Singapore, Singapore</p>
	<p><b>Radheshyam Chavan (Ph.d, M.Sc.)</b>  Associate Professor  Department of Toshniwal Arts, Commerce and Science College, Sengaon District Hingoli, Maharashtra, India  Email: rtcbotany@gmail.com  Phone: 08275949170</p>

	<p><b>Dr. P. Mehalingam (Ph.D)</b>  Assistant Professor  Department of Botany (DST-FIST Sponsored) V.H.N. Senthikumara Nadar College (An Autonomous Institution Affiliated to Madurai Kamaraj University, Madurai) Virudhunagar, Tamil Nadu, India  Email: mehalingamp@yahoo.co.in</p>
	<p><b>Kaptain Kishor Bajpayee (Ph.D.)</b>  Associate Professor  Center for Research in Ethno &amp; Medicobotany, Dr. R. M. L. Degree College (C.S.J.M. UNIVERSITY ) Allipur. Hardoi, India  Email: drkkb94@gmail.com</p>
	<p><b>Dr Jithesh M (PhD)</b>  Professor &amp; Head  Department of Kayachikitsa &amp; PG Studies in Manas roga Vaidyaratnam PS Varier Ayurveda College, Kottakkal Kerala, India  Email: drjitheshm@gmail.com</p>
	<p><b>Dr. Sr. M. Arul Sheeba Rani (PhD)</b>  Assistant Professor  Department of botany Nirmala College for Women Red Fields Coimbatore TamilNadu India  Email: sheebam582@yahoo.com  Phone: 9600775537</p>
	<p><b>Ramarajan S (M.Sc., M.Phil., Ph.D)</b>  Assistant Professor  Researcher Survey Medicinal Plant Unit [Central Council for Research in Siddha] Govt. Siddha Medical College Campus, Tamil Nadu, India  Email: sramabio@gmail.com  Phone: 9025935688</p>
	<p><b>Dr. Sapan Patel (Ph.D)</b>  Assistant Professor  School of Studies in Botany, Jiwaji University, Gwalior, Madhya Pradesh, India  Email: sapanpatel79@gmail.com  Phone: 9425068613</p>
	<p><b>Dr. Debabrata Das (Ph.D)</b>  Associate Professor  Department of Botany, Lalgarh Govt. College, Jhargram, West Bengal, India  Email: debabratlalgarh@gmail.com  Phone: 9800301026</p>
	<p><b>Dr. Praveen Dhar T (Ph.D. (Botany))</b>  Assistant Professor  Department of Botany, St. Stephen's College, Pathanapuram, Kollam, Kerala, India  Email: dharpraveent@gmail.com  Phone: +919446170833</p>
	<p><b>Suchita Sinha (Ph.D)</b>  Associate Professor and Head of the Department  Department of Botany, Lady Brabourne College, Kolkata, West Bengal, India  Email: suchita1sinha@yahoo.co.in  Phone: +919433122420</p>
	<p><b>Dr Janhavi Ashish Arekar (M. Sc. Botany, Ph. D. Botany)</b>  Associate Editor  Department of Botany Smt. Devkiba Mohansinhji College of Commerce and Science, Silvassa, Dadara and Nagar Haveli, India  Email: janhaviashisharekar@gmail.com  Phone: 7874182182</p>

	<p><b>Dr. M. Mathiyazhagan (M.Sc, Ph.D)</b> Associate Professor School of Basic &amp; Applied Sciences Apex Professional University Pasighat, Arunachal Pradesh, India Email: mathibdu@gmail.com Phone: 9751931230</p>
	<p><b>Dr. Aparna Amit Saraf (PhD (Plant Physiology and Biochemistry))</b> Associate Professor Department of Botany The Institute of Science, Mumbai, Maharashtra, India Email: draparnasaraf@yahoo.co.in Phone: 9869357636</p>
	<p><b>Hemantkumar Thakur (M.Sc. Ph.D.)</b> Assistant Professor Post Graduate &amp; Research Center of Botany Gokhale Education Society H.P.T. Arts &amp; R.Y.K. Science College Prin. T. A. Kulkarni Vidyanagar Nasik, Maharashtra India Email: hemant13570@gmail.com Phone: 9767199633</p>
	<p><b>Dr. Beena Kumari (M.Sc., Ph.D., LL.B.)</b> Associate Professor Department of Botany, Hindu College Moradabad (Affiliation MJP Rohilkhand University, Bareilly), Uttar Pradesh, India Email: beenakumari.botany@gmail.com Phone: 9412143577</p>
	<p><b>Dr. Alok Ranjan Sahu (M.Sc, M.Phil., Ph.D)</b> Principal Saraswati Degree Science College, Near Petrol Pump, Kesinga Road, Bhawanipatna, Odisha, India Email: alok.btgene@gmail.com Phone: +919692895348</p>
	<p><b>Dr. G. Manikandan (M.S., PhD)</b> Assistant Professor Department of Botany Sri Kaliswari College Sivakasi, Tamil Nadu, India Email: rgmani.19@gmail.com Phone: 9585164619</p>
	<p><b>Dr. Jeetendra Sainkhediya (PH.D.,M.Phil)</b> Assistant Professor Department of Botany Govt.PG College Sendhwa, Barwani Madhya Pradesh, India Email: jeetu.sainkhediya@gmail.com Phone: 9826613120</p>
	<p><b>Purshotam Singh (Ph.D)</b> Associate Professor Division of Agronomy, SKUAST-K, Shalimar, Srinagar, Kashmir, Jammu and Kashmir, India Email: drpurshotam@gmail.com Phone: 9419057547</p>
	<p><b>Dr. Lily Trivedi (Ph.D(BOTANY))</b> Associate Professor Jayoti Vidyapeeth Women's University, Vedaant Gyan Valley, Village-Jharna, Mahala Jobner Link Road, Jaipur Ajmer Express Way, Rajasthan INDIA Email: drlilytrivedi@jvwu.ac.in Phone: 9460587924</p>
	<p><b>Dr. Praveen Dhar T (M.Sc ( Botany) MPhil( Botany) MEd., MPhil( Education) M.A( English) MA ( History) Ph.D ( Botany))</b> Assistant Professor Department of Botany, St.Stephens College, Pathanapuram, Kollam Kerala, India Email: dharpraveent@yahoo.com Phone: 9446170833</p>

	<p><b>Anitha. T (M.Sc.,B.Ed.,Ph.D.,PGDCA)</b>  Assistant Professor  Department of Botany, Nirmala College for Women Coimbatore Tamilnadu India  Email: tanitha101982@yahoo.com  Phone: 919566915133</p>
	<p><b>Dr Sheeja T Tharakan (PhD)</b>  Assistant Professor  Department of Botany Vimala College (Autonomous) Thrissur Kerala India  Email: sheejatharakan@gmail.com  Phone: 9447991558</p>
	<p><b>Dr. Janhavi Ashish Arekar (M. Sc. Botany, Ph. D. Botany)</b>  Assistant Professor  Department of Botany, Smt. Devkiba Mohansinhji Chauhan College of Commerce and Science, Silvassa, Dadara and Nagar Haveli India  Email: janhaviashisharekar@gmail.com  Phone: 7874182182</p>
	<p><b>Dr. Shailesh Kumar (Ph.D.)</b>  Assistant Professor &amp; Scientist  Department of Botany &amp; Plant Physiology Faculty of Basic Sciences &amp; Humanities Dr.Rajendra Prasad Central Agricultural University, PUSA, Samastipur, Bihar, India  Email: shailesh_agri@yahoo.com  Phone: 9431013932</p>
	<p><b>Dr Praveen Dhar T (MSc(Botany) MPhil( Botany) MEd., MPhil(Education) MA. English., MA History., PhD(Botany))</b>  Assistant Professor  Department of Botany, StStephen's College, Pathanapuram., Kollam, Kerala, India  Email: dharpraveent@gmail.com  Phone: 9446170833</p>
	<p><b>Dr. Abhijeet Rajaram Kasarkar (MSc, PhD)</b>  Assistant professor  Department of Botany, Vivekanand College, Kolhapur, Maharashtra  Email: abhikasarkar@rediffmail.com  Phone: 9420494379</p>
	<p><b>Anand Anna Waghmare (M.Sc.,Ph.D.)</b>  Assistant professor  Toshniwal College, Department of Botany and Biotechnology At- Sengaon, Dist.-Hingoli Maharashtra, India  Email: anandwaghmare90@gmail.com  Phone: 7030245018</p>
	<p><b>Dr. Varsha Dinkarrao Hutke (M.Sc., Ph.D. Botany)</b>  Assistant Professor  Department of Botany, Government Vidarbha Institute of Science &amp; Humanities, Amravati, Maharashtra, India  Email: vdhutke@gmail.com  Phone: 09421789052</p>
	<p><b>Dr. Jai Knox (Ph.D)</b>  Assistant Professor  Department of Botany, Wilson College, Mumbai, Maharashtra, India  Email: drjaiknox@gmail.com  Phone: 9920081094</p>
	<p><b>Rp Singh (Ph.D)</b>  Assistant Professor  Department of Botany, DDU Gorakhpur University Gorakhpur, Uttar Pradesh, India  Email: ravinder.20033@gmail.com  Phone: 9811426055</p>


	<p><b>Ajeet Singh (Ph.D)</b>  Assistant Editor  Department of Botany and Microbiology, Gurukul Kangri University, Haridwar, Uttarakhand, India  Email: nikhilgupta321@gmail.com  Phone: 918791539165</p>
	<p><b>Dr. Sanjay Kumar Pandey (M.Sc., Ph.D)</b>  Assistant Professor of Botany  Department of Botany, S M P Government Degree College, Ballia, Uttar Pradesh, India  Email: drskp27@gmail.com  Phone: 09415837498</p>
	<p><b>Dr. Arijit Sinhababu (Ph.D.)</b>  Assistant Professor  Department of Botany, Bankura Christian College, Bankura, West Bengal, India  Email: arijitsinhababu@rediffmail.com</p>
	<p><b>T. Janaki (M.Sc., (Bot), M.Sc., (Micro), B.Ed., M.Phil, (Bot) Ph.D., (Bot))</b>  Assistant Professor  Department of Botany, KMCPGS, Lawspet, Puducherry, Tamil Nadu, India  Email: janakishaila@gmail.com</p>
	<p><b>Dr. Mangesh Jagatrao Dagawal (PhD)</b>  Head, Department of Botany  Smt. Radhabai Sarda Arts, Commerce &amp; Science College, Anjangoan Surji, Dist. Amravati (MS), India  Email: mdagawal@gmail.com  Phone: 09326827719</p>
	<p><b>Dr. Smruti Sohani (PhD)</b>  Associate Professor  Institute of Agriculture Science SAGE University, Indore (M.P), India  Email: smrutisohani@gmail.com  Phone: 7777880511</p>
	<p><b>Dr. Umesh Bhauraoji Kakde (PhD)</b>  Associate Professor  Botany at The Institute of Science, Mumbai, India  Email: drumeshkakde@gmail.com  Phone: 9869353455</p>
	<p><b>Dr. A. S. Jondhale (PhD)</b>  Head and Assistant Professor  M.J.M. Arts, Commerce and Science, College At/Po. Karanjali, Tal. Peth, Dist. Nashik, Maharashtra State, India  Email: avinashjondhale51@gmail.com  Phone: 08454845351</p>
	<p><b>Dr. Said Hassan (PhD)</b>  Assistant Professor  Bacha Khan College, Peshawar, Charsadda, Khyber Pakhtunkhwa, Pakistan  Email: saidhassan15@yahoo.com  Phone: +00923459151715</p>
	<p><b>Dr. Shahdab Hussain (PhD)</b>  Assistant Professor  Department of Botany, Sangam University, Bhilwara Rajasthan India  Email: drshahdabhussain@yahoo.com  Phone: 8233152670</p>

	<p><b>Dr. K. Rajkumar (PhD)</b>  Assistant Professor  Department of Botany, Laxminarayana Arts and Science College for Women, Dharmapuri, Periyar University, Salem, India.  Email: krajkumar2000@gmail.com  Phone: +918489570143</p>
	<p><b>Dr. Rania Ibrahim Mohammad Almoselhy (PhD)</b>  Associate Professor  Oils and Fats Laboratory, Food Technology Research Institute (FTRI), Agricultural Research Center (ARC) - Giza - Egypt  Email: rania_almoselhy@outlook.com  Phone: 002+01092497977</p>
	<p><b>Dr. V. Kadirvelmurugan (PhD)</b>  Assistant Professor  PG &amp; Research Department of Botany, Padmavani Arts and Science College for Women, Opp. Periyar University, Salem, Tamil Nadu, India  Email: v.kathirvelmurugan@gmail.com  Phone: +919488061705</p>
	<p><b>Dr. Jatinder Singh (PhD)</b>  Dean  Faculty of Agriculture, Sri Guru Granth Sahib World University, Fatehgarh Sahib, Punjab, India  Email: jatinder.19305@lpu.co.in  Phone: +919876268247</p>
	<p><b>Dr. Pranabesh Ghosh (PhD)</b>  Assistant Professor  School of Bio Science, Seacom Skills University, Santiniketan, Bolpur, Birbhum, West Bengal, India  Email: doctorpranabeshghosh@gmail.com  Phone: +919339522104</p>
	<p><b>Tamer Mansour (PhD)</b>  Assistant Professor  Department of Agricultural Economic, National Research Centre, Cairo, Egypt  Email: tamer_baz@yahoo.com</p>
	<p><b>Palvan Kalandarov</b>  Professor  Department of Automation and Control of Technological Processes, Tashkent Institute of Irrigation and Agricultural Mechanization Engineers, Tashkent, Uzbekistan  Email: eest_uz@mail.ru  Phone: +998935552552</p>
	<p><b>Rashmi Misra</b>  Assistant Professor  Food, Nutrition and Dietetics (FND) Department, Faculty of Agriculture (FOAG) in Sri Sri University, Cuttack, Odisha, India  Email: rasmi.minu1@gmail.com  Phone: 8249832027</p>
	<p><b>Anupam Dakua (PhD)</b>  Assistant Professor  Department of Extension Education, Faculty of Agriculture (FOAG), Sri Sri University, Cuttack, Odisha, India  Email: anupamdakua2@gmail.com  Phone: 7309665143</p>
	<p><b>M. Cuneyt Bagdatli (PhD)</b>  Associate Professor  Department of Biosystem Engineering, Faculty of Engineering and Architecture, Nevsehir Haci Bektas Veli University, Nevsehir, Turkey  Email: cuneytbagdatli@gmail.com  Phone: +903842281000</p>

	<p><b>Lonel Bondoc (PhD)</b>  Associate Professor  Department of Public Health, Faculty of Veterinary Medicine Iasi, University of Agricultural Sciences and Veterinary Medicine Iasi, Romania  Email: bondoc_mv@yahoo.com</p>
	<p><b>Dr. Abdelhak Rhouma</b>  Lecture  Department of Plant Protection and Environmental Health (Phytopathology), Higher Agronomic Institute of Chott Mariem Sousse, University of Sousse, Tunisia  Email: rhouma.abdelhak@gmail.com  Phone: 21650829121</p>
	<p><b>Darwin H Pangaribuan (PhD)</b>  Associate Professor  Faculty of Agriculture, University of Lampung, Indonesia  Email: darwin.pangaribuan@fp.unila.ac.id  Phone: +6282183283777</p>
	<p><b>Dr. Mohammad Asadi</b>  Assistant Professor  Department of Plant Protection, University of Mohaghegh Ardabili, Ardabil, Iran  Email: assadi20@gmail.com</p>
	<p><b>Dr. S. Dineshkumar</b>  Assistant Professor  Department of Agricultural Extension, Thanthai Roever Institute of Agriculture and Rural Development, Valikandapuram, Perambalur District, Tamil Nadu, India  Email: dinezselsva@gmail.com  Phone: +91 6380272599</p>
	<p><b>Dr. P. Jaisankar</b>  Assistant Professor  Department of Horticulture, Thanthai Roever Institute of Agriculture and Rural Development (TRIARD), Perambalur, Tamil Nadu, India  Email: sankarjai03@gmail.com  Phone: 9688860521</p>
	<p><b>R. Karthikeyan (M.Pharm., Ph.D.)</b>  Associate Professor  Department of Pharmacognosy (Medicinal Botany), Vignan Pharmacy College, Vadlamudi, Andhra Pradesh, India  Email: rkcognosy@gmail.com  Phone: 9966847127</p>
	<p><b>Dr. Khongdet Phasinam</b>  Assistant Professor  Department of Faculty of Food and Agricultural Technology, Pibulsongkram Rajabhat University, Thailand  Email: phasinam@psru.ac.th</p>
<p><b>Associate Editors</b></p>	
	<p><b>Dr. Souvik Tewari (M.Sc., Ph.D.)</b>  Assistant Professor  Department of Food and Nutrition, Swami Vivekananda University, Barrackpore, West Bengal, India  Email: souviktewari@gmail.com  Phone: 9002435977  ORCID ID: <a href="https://orcid.org/0000-0002-1463-2290">https://orcid.org/0000-0002-1463-2290</a></p>

## Vol. 5, Issue 1, Part B (2023)

S. No.	Title and Authors Name
1	<p><b>Precision agriculture: Disruption and transformation of the agriculture and food industry</b> Charles McMillan DOI: <a href="https://doi.org/10.33545/2664844X.2023.v5.i1b.127">https://doi.org/10.33545/2664844X.2023.v5.i1b.127</a> Int. J. Agric. Food Sci., 2023; 5(1): 101-106 <a href="#">Abstract</a>   <a href="#">Download</a>   Country: Canada   File Size: (775 KB)   60 Views 17 Downloads</p>
2	<p><b>Social economic factors influencing adoption of non-farm poverty coping strategies in Kuje area council, Abuja, Nigeria</b> Callister Amina Ekele, Samson Olayemi Sennuga, Joseph Bamidele, Funso Omolayo Alabuja and Bankole Osho-Lagunju DOI: <a href="https://doi.org/10.33545/2664844X.2023.v5.i1b.129">https://doi.org/10.33545/2664844X.2023.v5.i1b.129</a> Int. J. Agric. Food Sci., 2023; 5(1): 107-116 <a href="#">Abstract</a>   <a href="#">Download</a>   Country: Nigeria   File Size: (733 KB)   55 Views 15 Downloads</p>
3	<p><b>Insects pest repellent, essential oils, is can be an efficacious alternative to synthetic pesticides</b> Mubarak Abdelrahman Salim Eisa, Olha Matsera and Ludovít Cagáň DOI: <a href="https://doi.org/10.33545/2664844X.2023.v5.i1b.130">https://doi.org/10.33545/2664844X.2023.v5.i1b.130</a> Int. J. Agric. Food Sci., 2023; 5(1): 117-125 <a href="#">Abstract</a>   <a href="#">Download</a>   Country: Slovakia   File Size: (1067 KB)   71 Views 32 Downloads</p>
4	<p><b>Influence of electrical process parameters to inactivate microorganisms in orange juice by PEF technology</b> S Krishnaveni DOI: <a href="https://doi.org/10.33545/2664844X.2023.v5.i1b.131">https://doi.org/10.33545/2664844X.2023.v5.i1b.131</a> Int. J. Agric. Food Sci., 2023; 5(1): 126-132 <a href="#">Abstract</a>   <a href="#">Download</a>   Country: India   File Size: (1052 KB)   52 Views 26 Downloads</p>
5	<p><b>India's agriculture: Issues and priorities</b> Dr. Nazeerudin and Dr. BM Rajani Kanth DOI: <a href="https://doi.org/10.33545/2664844X.2023.v5.i1b.132">https://doi.org/10.33545/2664844X.2023.v5.i1b.132</a> Int. J. Agric. Food Sci., 2023; 5(1): 133-136 <a href="#">Abstract</a>   <a href="#">Download</a>   Country: India   File Size: (720 KB)   39 Views 10 Downloads</p>
6	<p><b>Assessment of climate variability and effective coping strategies used by rice farmers in Abuja, Nigeria</b> Cecilia Ijachi, Samson Olayemi Sennuga, Osho-Lagunju Bankole, Emeka Franklin Okpala and Timipre Joseph Preyor DOI: <a href="https://doi.org/10.33545/2664844X.2023.v5.i1b.133">https://doi.org/10.33545/2664844X.2023.v5.i1b.133</a> Int. J. Agric. Food Sci., 2023; 5(1): 137-143 <a href="#">Abstract</a>   <a href="#">Download</a>   Country: Nigeria   File Size: (766 KB)   46 Views 22 Downloads</p>
7	<p><b>Effect of diet on human gut health: A review</b> Payel Kumar Roy, Anup Ghosal, Titlee Majumder and Swastika Pradhan DOI: <a href="https://doi.org/10.33545/2664844X.2023.v5.i1b.134">https://doi.org/10.33545/2664844X.2023.v5.i1b.134</a> Int. J. Agric. Food Sci., 2023; 5(1): 144-146 <a href="#">Abstract</a>   <a href="#">Download</a>   Country: India   File Size: (593 KB)   180 Views 39 Downloads</p>
8	<p><b>Recent developments in ready-to-eat and ready-to-cook foods: An overview</b> Ushas Mathew and Dr. Priyanka Sharma DOI: <a href="https://doi.org/10.33545/2664844X.2023.v5.i1b.135">https://doi.org/10.33545/2664844X.2023.v5.i1b.135</a> Int. J. Agric. Food Sci., 2023; 5(1): 147-152 <a href="#">Abstract</a>   <a href="#">Download</a>   Country: India   File Size: (637 KB)   28 Views 11 Downloads</p>



ISSN Print: 2664-844X  
 ISSN Online: 2664-8458  
 Impact Factor: RJIF 5.6  
 IJAFS 2023; 5(1): 117-125  
[www.agriculturaljournals.com](http://www.agriculturaljournals.com)  
 Received: 06-10-2022  
 Accepted: 13-11-2022

**Mubarak Abdelrahman Salim Eisa**

<sup>1</sup> Institute of Agronomical Sciences, Faculty of Agrobiological and Food Resources, Slovak University of Agriculture Nitra, Tr. A. Hlinku 2, 949 76, Nitra, Slovakia

<sup>2</sup> Department of Plant Protection, University of Zalingei in Sudan, Faculty of Agriculture, Sudan

**Olha Matsera**

Institute of Agronomical Sciences, Faculty of Agrobiological and Food Resources, Slovak University of Agriculture Nitra, Tr. A. Hlinku 2, 949 76, Nitra, Slovakia

**Ludovít Cagaň**

Institute of Agronomical Sciences, Faculty of Agrobiological and Food Resources, Slovak University of Agriculture Nitra, Tr. A. Hlinku 2, 949 76, Nitra, Slovakia

**Corresponding Author:**  
**Mubarak Abdelrahman Salim Eisa**

<sup>1</sup> Institute of Agronomical Sciences, Faculty of Agrobiological and Food Resources, Slovak University of Agriculture Nitra, Tr. A. Hlinku 2, 949 76, Nitra, Slovakia

<sup>2</sup> Department of Plant Protection, University of Zalingei in Sudan, Faculty of Agriculture, Sudan

## Insects pest repellent, essential oils, is can be an efficacious alternative to synthetic pesticides

**Mubarak Abdelrahman Salim Eisa, Olha Matsera and Ludovít Cagaň**

DOI: <https://doi.org/10.33545/2664844X.2023.v5.i1b.130>

### Abstract

The aim of this study was to verify the degree of repellent effect of two chemical substances (DEET and 2-undecanone) and three essential oils (EOs), *Allium sativum*, *Lavandula angustifolia* and *Eucalyptus globulus*. *Leptinotarsa decemlineata* Say. Were exposed to different concentrations (0.2, 0.4 and 0.6%) in a two-choice olfactometer and (0.02, 0.04 and 0.06 ml/cm<sup>2</sup>) area preference assay. The effect was recorded 15 minutes, in olfactometer and 6-24 hours in area preference test, after the application of the treatments. It was found that *A. sativum*, *E. globulus*, and *L. angustifolia* had a significant repellent effect. We also found that the repellent effect increased with increasing concentration and time. An olfactometer and area preference assay achieved the highest repellency index (%) from (*A. sativum*, *E. globulus*, *L. angustifolia*, deet, and 2-undecanone), indicating RI 80% (15mins) and RI 93% at 6-24 hours in concentration (0.6%, 0.06 ml/cm<sup>2</sup>). The present results show that the tested oils have high activity against *L. decemlineata*. We concluded that the EOs may have potential as an alternative to chemical control, all these efficient EOs could be used as effective biocontrol agents against various field pests.

**Keywords:** Repellent, olfactometer, essential oils, synthetic pesticides, *L. decemlineata*

### Introduction

The Colorado potato beetle (*Leptinotarsa decemlineata* Say) is the most important pest of potato (FAO STAT, 2020) <sup>[9]</sup>. It is widespread in North America, Europe, and Asia, covering about 16 million km<sup>2</sup>, and continues to spread (James, 2011) <sup>[20]</sup>. In the 20th century, this pest became a major problem throughout Europe, Asia, and China (Oerke, 2006) <sup>[27]</sup>. Defoliation can reduce tuber yield by, more than 50% (James, 2011) <sup>[20]</sup>. The first recorded case of plant protection by chemical insecticide was found in the 1950s after the application of DDT (James 2011) <sup>[20]</sup>. Over time, CPB developed resistance to many insecticides which were initially effective (FAO STAT, 2020) <sup>[9]</sup>. Up to day, more than 300 cases of *L. decemlineata* resistance to 56 insecticidal agents have been detected (FAO STAT, 2020) <sup>[9]</sup>. In addition, different alternative chemical control methods have been introduced since the 19th century that have minimized CPB infestations. Crop rotation, trap crops, and other agrotechnical practices have been recommended to farmers (Hin, 2009) <sup>[14]</sup>. It has been concluded that the only two reliable methods are hand picking and mating with green. Hand picking, especially before mating, was found to be very effective but impractical on a larger scale. Therefore, most alternative methods for CPB management have failed, and it's high time to look for more economical and sustainable means (Isman, 2006) <sup>[15]</sup>.

EOs can contain many terpenes and low molecular weight phenolic substances and are easily extracted from plant material by steam distillation (Regnault-Roger, 2008) <sup>[31]</sup>. Studies in several countries have confirmed that some plant essential oils repel insect pests and used insecticidal activity on certain pests upon contact or by fumigation (Regnault-Roger, 2008 & Mukarram *et al.*, 2021a) <sup>[31, 26]</sup>. Several commercial bioinsecticides are based on plant extracts (essential oils) or their chemical compositions in many major agricultural regions, including: Europe, USA, China, and Latin America (Isman, 2017) <sup>[16]</sup>. The study literature reports that the plant species with insect repellent activities and their active ingredients have been published by (Isman & Machal 2006; Khallaayoune *et al.*, 2009; González-Coloma *et al.*, 2010 & Regnault-Roger 2013) <sup>[18, 22, 12, 32]</sup>, includes *Aartemisia vulgaris* L. (Thujone,

cineole), *Cinnamomum camphora* (L.) Presl (cinnamaldehyde), *Curcuma longa* L. (turmerone), *Eucalyptus globulus* Labill, *Myrtus communis* L. and *Rosmarinus officinalis* L. (cineoles), *Juniperus virginiana* L. ( $\alpha$  and  $\beta$  pinene, methyl-eugenol), *Lavandula angustifolia* Mill, (linalool, linalyl acetate), *Litsea cubeta* Pers., and *Cymbopogon* species (*citral*, *citronellal*, *citronellol*), *Melaleuca leucadendron* L. (terpineol,  $\gamma$ -terpinene), *Mentha pulegium* L. (pulegone), *Mentha piperita* L. (menthone, menthol), *Nepeta cataria* L. (nepetalactone), *Pelargonium* sp. (Geraniol), *Syzygium aromaticum* L. (eugenol), *Thymus* sp. And *Origanum vulgare* L. (thymol, carvacrol, p-cymene).

In a laboratory investigation, the essential oils of *Allium sativum* Linnaeus, 1793 showed insecticidal activities on different Coleoptera pests (Abdalla *et al.*, 2017) [1]. Garlic extract had effective repellent activity against *Tribolium castaneum* (Herbst, 1797) (Jahromi *et al.*, 2012) [19]. The major chemical compositions of *eucalyptus* oils are 1, 8-cineole and  $\alpha$ -pinene or 1, 8-cineole and linalool (Boukhatem *et al.*, 2020) [3]. The essential oil of *Allium sativum* Linnaeus, 1793, comprise sulphur-containing organic molecules, mainly diallyl disulfide, allyl propyl disulfide, allyl sulfoxide, and allicin. The activity of diallyl disulfide as a fumigant against many insects and pathogens was qualified by Chiam *et al.*, (1999) [5]. Various essential oils are used in registered commercial biopesticides, especially in developed countries. Among these products, garlic oil is the most commonly used (Regnault *et al.*, 2012) [30]. The most effective and well-known insect repellents currently on the market are DEET (N, N-diethyl-3-methylbenzamide) and 2-undecanone (Maia, & Moore, 2011) [24]. DEET, an active ingredient used worldwide since 1946, is a highly effective repellent for a variety of insect species (Maia, & Moore, 2011) [24]. 2-undecanone (methyl nonyl ketone), a natural non-toxic insect repellent compound, has been found in palm kernel oil and soybean oil (Sanghong *et al.*, 2015) [33].

Considering the needs for sustainable and environmentally friendly pesticides and the efficacy of EOs. The aim of our study was to investigate the repellency of *A. sativum*, *L. angustifolia* and *E. globulus* EO as plant secondary metabolites against Colorado potato beetle under laboratory conditions.

## Materials and Methods

### Basic experiment information

This study was carried out at the Laboratory of Entomology, Faculty of Agrobiological and Food Resources, Slovak University of Agriculture in Nitra. Laboratory investigations were divided into two experiments. The experiment included "two-choice olfactometer" repellency and Area preference assay with eight EOs, two chemicals against the *L. decemlineata*. The study was carried out under the laboratory conditions at the temperature of  $20 \pm 1$  °C, humidity  $50 \pm 1$  %, and /light/dark 16:08/ h.

**Insects used:** Adults of the *L. decemlineata* were collected from a potato culture at a local field in Nitra (Slovakia) 48°18'16.7"N 18°05'41.3"E in August-September 2021. Nutrients were potato tubers as adult foods. Approximately 1000 adults were incubated at  $20 \pm 1$  °C and 50% relative humidity (RH) under a long day (16/8 h) photoperiod in plastic boxes (74 cm x 52 cm x 28 cm). Adults of both sexes of the approximately same age were used in the experiments.

### Essential oils and chemical substances used

Essential oils were from plant species *A. sativum*, *L. angustifolia* and *E. globulus*. The most important chemicals in *A. sativum* EO were diallyl disulphide, diallyl trisulphide, and diallyl sulphide, in the case of *L. angustifolia* they were citral, coumarin, eugenol, geraniol, limonene, linalool, in EO from *E. globulus* they were 1,8 cineole, dipentene, alpha phellandrene, beta pinene. Commercial chemical substances were purchased from Sigma-Aldrich (DEET (N, N-diethyl-3-methylbenzamide) at concentration 97%. 2-Undecanone (methyl nonyl ketone) in concentration 99%. According to the website of this company and Fusková, & Cagaň, (2021) [10].

([https://www.mysticmomentsuk.com/collections/essential-oils?grid\\_list=grid-view](https://www.mysticmomentsuk.com/collections/essential-oils?grid_list=grid-view)).

**Table 1:** Dosage used in laboratory experiment

EOs/Dose ( $\mu$ l)	Dose (ml)	Solvent (acetone)/ ml	Concentration%
20	0.02	10	0.2
40	0.04	10	0.4
60	0.06	10	0.6

$$V/V\% = [(Volume\ of\ solute) / (Volume\ of\ solution)] \times 100\%$$

### Application of essential oils in two-choice olfactometer test

The experiments were conducted using a Y-tube-olfactometer. A 0.2%, 0.4% and 0.6% tested essential oils of each solution was dosed on a piece of filter paper (No. 2, 10x20mm), odour sources were placed in the treatment arm, and a filter paper treated with 10  $\mu$ l of acetone was placed in the control side. Five replicates were used for each concentration%. Each replicate consisted of 5 responding CPB adults were leased individually into the olfactometer using flexible forceps. The measurement started after 15 minutes. The choices of the CPB were recorded. The calculated repellency index was % RI =  $(1 - (\%T / \%C)) \times 100$ , where % T is the percentage of insects on the treated arm and % C is the percentage of individuals on the control arm.

### Area preference assay

The repulsive effect was determined by the area preference method on filter paper as described by (Abdel-Tawab, 2016) [2]. Filter paper (Whatman N°1) with a diameter of 9 cm was cut into two halves, and 10  $\mu$ l of each concentration (0.02, 0.04, and 0.06 ml/cm<sup>2</sup>) of treated Eos was applied separately to one half of the filter paper using a pipette. The second half (control) was treated with 10  $\mu$ l of the solvent (acetone).

Both halves of filter paper were allowed to dry at room temperature for 10 minutes. The test insects (five adults) were released in the center of each filter paper and the Petri dish was immediately covered. Each treatment was repeated three times. The average duration of a repetition was 6 and 24 hours. The coverings of the Petri dishes were ventilated through tiny-holes whose diameter was tiny-enough to

prevent insects from escaping. Petri dishes were labeled to prevent mixing of treated and untreated sides. The repellency percent (PR%) of the treatment oils was calculated using the following formula: The calculated repellency index was  $\% RI = (1 - (\%T / \%C)) \times 100$ , where %T is the percentage of insects on the treated arm and % C is the percentage of individuals on the control arm.

**Table 2:** The scheme of the experiment №.

Factor A: Concentration,% and ml/cm <sup>2</sup>		Factor B: Time after treatment				Factor C – EOs for treatments	
1.	0.2	Olfactometer		Area preference		1.	<i>E. globulus</i>
2.	0.4	Minutes		Hours		2.	<i>L. angustifolia</i>
3.	0.6	1.	15	1.	6	3.	<i>A. sativum</i>
//	//	//	//	2.	24	4.	<i>Deet</i>
//	//	//	//	//	//	5.	<i>2-Undecanone</i>

**Table 3:** Chemical constituents (%) of *A. sativum*, *E. globulus*, and *L. angustifolia* essential oils included in repellency bioassays.

Compounds	Relative amount%		
	<i>A. sativum</i>	<i>E. globulus</i>	<i>L. angustifolia</i>
Limonene	-	≤28.75	≤0.74
Linalool	-	≤1.66	≤35.75
Citral	-	≤28.53	≤0.04
Eugenol	-	≤0.14	≤0.00
Methyl allyl sulfide	≤2.39	-	-
Dimethyl disulfide	≤0.543	-	-
Diallyl sulfide	≤5.283	-	-
Methyl allyl disulfide	≤11.293	-	-
Dimethyl trisulfide	≤0.628	-	-
Diallyl disulfide	≤37.231	-	-
Methyl allyl trisulfide	≤8.338	-	-
Diallyl trisulfide	≤18.287	-	-
Diallyl tetrasulfide	≤3.819	-	-
Geraniol	-	≤6.43	≤0.66
Citronellol	-	≤0.67	≤0.00

According to Fusková, & Cagaň, (2021) <sup>[10]</sup> and the web page of this company ([https://www.mysticmomentsuk.com/collections/essential-oils?grid\\_list=grid-view](https://www.mysticmomentsuk.com/collections/essential-oils?grid_list=grid-view))

## Results


**Two-choice olfactometer test:** All of the investigated plant essential oils *A. sativum*, *E. globulus*, *L. angustifolia* essential oil and two Deet and 2-Undecanone against *L. decemlineata* at all higher of the tested concentrations 0.6% caused a repellency index% higher a 80% (Table 5). On all 15 minutes of the investigation, the highest RI% rate of the *L. angustifolia* adults was observed at the highest concentration 0.6% with an *A. sativum*, *L. angustifolia* and

2-Undecanone. On the other hand, the lowest concentration 0.2% was observed the lowest repellency index 20% at all treated substances. The average repellency activity were increases with increasing the concentration and time 0.2% (0.7), 0.4% (0.5) and 0.6% (0.3) (Table 4). The average repellency activity was observed at the all concentrations and times, it was as follows: A “2-Undecanone (0.2), *A. sativum* (0.3), *L. angustifolia* (0.3), Deet (0.3) and *E. globulus* (0.4)” (Table 4).

**Table 4:** The average amount of *L. decemlineata* through experiment depending on the 1) concentrations%, (2), time per minutes (3), treated EOs and chemicals after application and (4) overall average number of insects in treated side.


(1) Concentration %	(4) Average number of insects/concentration	(2) Time after treatment, min	(4) Average number of insects/min	(3) EOs and chemicals	(4) Average number of insects/ EOs and chemicals
0.2	0.7				
0.4	0.5	15	1.5	<i>A. sativum</i>	0.3
0.6	0.3	//	//	<i>E. globulus</i>	0.4
//	//	//	//	<i>L. angustifolia</i>	0.3
//	//	//	//	DEET	0.3
//	//	//	//	2-Undecanone	0.2

Concentrations %, (2), time per minutes (3), treated EOs and chemicals after application and (4) overall average number of insects in treated side.



Source: made by authors based on the own research

Fig 1: The correlation field between average amount of *L. decemlineata* (X) and concentration levels of all EOs (Y).



Source: made by authors based on the own research

Fig 2: The correlation field between average amount of *L. decemlineata* (X) and overall treated EOs (Y).

**Table 5:** The average repellency effect of the chemicals and EOs to the adults of Colorado potato beetle, *Leptinotarsa decemlineata*. The repellency was based on two-choice olfactometer test. Five adults were included to each two-choice olfactometer test. Five replications were done for each concentrations (0.02, 0.04 and 0.06%). For each replication duration was 15 minutes. Standard deviation SD ±, Data variation,% and the calculated repellency index was  $\% RI = (1 - (\%T/\%C)) \times 100$ , where % T is the percentage of insects on the treated arm and % C is the percentage of individuals on the control arm.

Concentration,%	Time after, min	Variation indexes		Data variation,%	Control	RI,%
		Average	Standard deviation, SD			
<i>A. sativum</i>						
0.2	15	0.8	0.5	1.3	1	20
0.4	15	0.4	0.5	2.5	3	60
0.6	15	0.2	0.4	4.3	4	80
<i>E. globulus</i>						
0.2	15	0.8	0.7	1.8	1	20
0.4	15	0.8	1.0	2.5	1	20
0.6	15	0.6	0.4	1.5	2	40
<i>L. angustifolia</i>						
0.2	15	0.8	0.1	0.2	1	20
0.4	15	0.4	0.5	2.5	3	60
0.6	15	0.2	0.4	4.3	4	80
DEET						
0.2	15	0.6	0.8	2.8	2	40
0.4	15	0.4	0.2	0.9	3	60
0.6	15	0.4	0.5	2.5	3	60
2-Undecanone						
0.2	15	0.6	0.8	2.8	2	40
0.4	15	0.4	0.9	4.3	3	60
0.6	15	0.2	0.4	4.3	4	80

**Results of Repellency activity of an area preference assay**


The results of repellent activity of *A. sativum*, *E. globulus*, *L. angustifolia* (EOs) and two chemical substances deet and 2-undecanone against *L. decemlineata* are summarized in (Table 7 and 8), respectively. All essential oil and the chemicals showed significant repellent activity (RI%) at the concentrations used. An average number of repellency effects was observed between treated and untreated half filter paper disk for the oil formulation (0.2, 0.4, and 0.6 ml/cm<sup>2</sup>). The repellency index (RI%) was range 73 to 93% after 6 hours and 80 to 93% after 24 hours for all EOs (*A. sativum*, *E. globulus*, *L. angustifolia*, deet and 2-undecanone) and concentrations (0.2, 0.4 and 0.6 ml/cm<sup>2</sup>). (table 6) shows the average number of insects on the treated

side, all essential oils showed mode rate repellency activity lower with an average repellency values of 2.8 at 0.02 µl/cm<sup>2</sup>. The repellency was strongest in the first 24 hours after treatment. This pattern was observed in Table 7 and 8, which show that Eos achieved the highest repellency index (%) using CPB from (*A. sativum*, *E. globulus*, *L. angustifolia*, deet, and 2-undecanone), indicating 93% repellency at 6 and 24 hours in concentration (0.06 ml/cm<sup>2</sup>). The average repellency activity increased with increasing concentration and time 0.2 (2.8), 0.4 (1.9) and 0.6 ml/cm<sup>2</sup> (1.7), (table 6). The average repellency activity was observed at all concentrations and times, and was as follows: A "2-undecanon (0.5), *A. sativum* (0.5), *L. angustifolia* (0.6), *E. globulus* (0.9), and Deet (1.0)" (table 6).

**Table 6:** The average amount of *L. decemlineata* through experiment depending on the (1) concentrations ml/cm<sup>2</sup>, (2), time per hours (3), treated EOs and chemicals after application and (4) overall average number of insects in treated side.


(1) Concentration ml/cm <sup>2</sup>	(4) Average number of insects	(2) Time, hours	Average number of insects per (concentrations and hours)	(4) Average number of insects total time	(3) Eos and chemicals	(4) Average number of insects
0.2	2.8	6	1.6	3.7	<i>A. sativum</i>	0.5
0.4	1.9		1.1 1.0		<i>E. globulus</i>	0.9
0.6	1.7		1.2		<i>L. angustifolia</i>	0.6
//	//	24	0.8	2.7	DEET	1.0
//	//		0.7		2-Undecanone	0.5
//	//					

Concentrations ml/cm<sup>2</sup>, (2), time per hours (3), treated EOs and chemicals after application and (4) overall average number of insects in treated side.



Source: made by authors based on the own research

**Fig 3:** The correlation field between average amount of *L. decemlineata* (X) and concentration levels of all EOs (Y).



Source: made by authors based on the own research

**Fig 4:** The correlation field between average amount of *L. decemlineata* (X) and overall treated EOs (Y).

**Table 7:** The average repellent efficacy of the chemicals and of Eos against adults *L. decemlineata* after 6 hours. Repellent efficacy was based on area preference test. Three replicates were used for each concentration (0.02, 0.04, and 0.06 ml/cm<sup>2</sup>). Standard deviation SD ±, data

variation,%, and Repellency index was calculated as  $\%RI = ((1 - (\%T/\%C)) \times 100$ , where %T is the percentage of beetles on the treated filter paper half-disk and %C is the percentage of individuals on the control filter paper half-disk in the Petri dish. Five adults were placed in each Petri dish.

Repellent	Concentrations/ ml/cm <sup>2</sup>	Average number of adults counted after 6 hours				
		Average	SD	Control	Data variation,%	RI,%
<i>A. sativum</i>	0.2	0.7	0.6	4.3	8.7	87
	0.4	0.3	0.6	4.7	17.3	93
	0.6	0.3	0.6	4.7	17.3	93
<i>E. globulus</i>	0.2	1.3	0.6	3.7	4.3	73
	0.4	0.3	0.6	4.7	17.3	93
	0.6	1.3	2.3	3.7	17.3	73
<i>L. angustifolia</i>	0.2	0.7	1.2	4.3	17.3	87
	0.4	0.7	1.2	4.3	17.3	87
	0.6	0.7	1.2	4.3	17.3	87
DEET	0.2	1.3	1.2	3.7	8.7	73
	0.4	1.0	1.0	4.0	10.0	80
	0.6	0.7	1.2	4.3	17.3	87
2-Undecanone	0.2	0.3	0.6	4.7	17.3	93
	0.4	0.7	0.6	4.3	8.7	87
	0.6	0.3	0.6	4.7	17.3	93

**Table 8:** The average repellency effect of the chemicals and Eos against adults of *L. decemlineata* after 6 hours. The repellency was based on area preference assay. Three replications were used for each concentrations (0.02, 0.04 and 0.06 ml/cm<sup>2</sup>). Standard deviation SD  $\pm$ , Data variation,% and The repellency index was calculated as  $\%RI = ((1 - (\%T/\%C)) \times 100$  where % T is the percentage of beetles on the treated half-disk of filter paper and % C is the percentage of individuals on the control half-disk of filter paper in the Petri dish. Five adults were included in each Petri dish.

Repellent	Concentrations/ ml/cm <sup>2</sup>	Average number of adults counted after 24 hours				
		Average	SD	Control	Data variation,%	RI,%
<i>A. sativum</i>	0.2	0.3	0.6	4.7	17.3	93
	0.4	0.3	0.6	4.7	17.3	93
	0.6	0.3	0.6	4.7	17.3	93
<i>E. globulus</i>	0.2	1.0	1.0	4.0	10.0	80
	0.4	0.3	0.6	4.7	17.3	93
	0.6	0.3	0.6	4.7	17.3	93
<i>L. angustifolia</i>	0.2	0.3	0.6	4.7	17.3	93
	0.4	0.3	0.6	4.7	17.3	93
	0.6	0.3	0.6	4.7	17.3	93
DEET	0.2	1.0	1.0	4.0	10.0	80
	0.4	0.7	0.6	4.3	8.7	87
	0.6	0.3	0.6	4.7	17.3	93
2-Undecanone	0.2	0.7	0.6	4.3	8.7	87
	0.4	0.3	0.6	4.7	17.3	93
	0.6	0.3	0.6	4.7	17.3	93

## Discussion

Insect repellents are substances that extracted locally to keep insects from flying onto, settling on, or stinging the skin of human or animals. Certain volatile plant compounds and essential oils derived from plants are known to repel various pest species and are considered minimal risk pesticides. The repellency of the essential oils and the two chemicals against adults *L. decemlineata* is shown in Tables 5, 7, and 8, where the repellency generally increased with increasing concentration. On the other hand, the lowest repellency was observed at the lowest concentrations for all concentrations tested. The highest repellency (93%) of the oils were found at the highest concentrations with the area preference-based test, and the highest repellency rate (80%) of the two-choice odour test was also found at the same highest concentrations. Some study was found the fruits of *Melia azedarach* showed excellent repellency against many insects as reported by Panji, (1964) [28], who found that a 5% ethanolic extract of *Melia azedarach* repelled adults *Aulacophora foveicollis* L. Kebede *et al.*, (2010) [21], found that a 5% ethanolic extract of *Melia azedarach* repelled at least 30% of *Raphidopalpa foveicollis* Lucas beetles (1

hour) and a maximum of 65% of beetles (48 hours), while a 10% extract repelled a maximum of 76% of beetles in 48 hours. *Melia azedarach* oil at 2% concentration provided 95.13%, for 7 hours and 20 minutes, while the 5% oil provided 96.20%, for 8 hours and 20 minutes against *Phlebotomus orientalis* (vector of visceral leishmaniasis), (Kebede *et al.*, 2010) [21]. Our study showed the potential of DEET and 2-undecanone and especially *A. sativum*, *E. globulus*, *L. angustifolia* essential oil against *L. decemlineata*. In last years, the number of research publications on the use of plant biopesticides as pest control agents in integrated pest management has increased exponentially. In addition, many laboratories worldwide have concentrated studies on essential oils and their constituents to exploit their biopesticides properties (Grumezescu, 2017) [13]. However, despite these advances, there are still few commercially available products based on essential plant oils or their isolated components (Dougoud *et al.*, 2019) [8]. Recently, the use of repellents has attracted the interest of scientists and the crop protection industry (Isman, 2000) [17]. The most promising are DEET and 2-undecanone,

which are already used as ingredients in insect repellents (Witting-Bissinger *et al.*, 2008) [36].

Extracts of *Allium sativum* have considerable acaricidal and insecticidal properties against Coleoptera and Diptera pests (Abdalla *et al.*, 2017) [1]. As well as the results in (Plata-Rueda *et al.*, 2017 & Moncada *et al.*, 2021) [29,25]. As reported by Fusková, & Cagán, (2021) [10], who found that the high repellency (90%) against *Tenebrio molitor* and high affinity to *A. obtectus* and *T. confusum*. Strikingly, high efficacy (95%) was observed even at the lowest concentration (0.01%). Our results confirmed high repellency (93% within 6 and 24 hours) against *L. decemlineata*, with the highest concentration (0.06%) based on an area preference test and a two-choice odour test.

The use of *Eucalyptus* EO in our study showed the results of high repellency (93% within 24 hours), with the odor test based on area preference test and two choice, was found in the highest concentration (0.06%). Similar results were also obtained by (Shahriari *et al.* 2019) [34]. *Eucalyptus globulus* EO had a significant effect on *T. confusum* and *A. obtectus*, which was similar to the repellency previously determined for *E. kuehniella* (80% within 24 hours).

*L. angustifolia* EO confirmed in our results a high repellency (93% within 6 and 24 hours) against *L. decemlineata*, with the highest concentration (0.06%) based on an area preference test and a two-choice odor test. According to the research of Germinara *et al.*, (2017) [11], the repellent activity of *L. angustifolia* EO seems to be high against all the pests tested (80%). The effect was detectable even at low concentrations, which is consistent with the results obtained for *S. granaries*. Moreover, the repellent effect was also significant, as also confirmed in various studies with different EOs (Chaudhari *et al.*, 2021; Tu *et al.*, 2017 & Conboy *et al.*, 2020) [4,35,6]. Thus, we can conclude that these EOs may have significant potential against a wide range of pests.

In summary, chemicals pesticides and repellency their kill rapidly target pests and provide highly control when applied. However, their application leaves toxic residues in field crops, soil, and stored grains that are harmful to consumers. Therefore, a new and valuable alternative approach should be found to reduce the use of insecticides in field warehouses and stored grains. One possible method is the encapsulation and emulsification of plant EOs as surfactants in greenhouses (De Oliveira *et al.*, 2019) [7]. The second is the direct application of EOs. Both approaches could be powerful tools for integrated management of field and storage pests. However, more research needs to be done in the laboratory and in the field.

## Conclusions

Novel insect pests repellency of the essential oils in our study was not established on particularly chemicals compositions to essential oils. The tested plant essential oils had the negative impact on *L. decemlineata*. Our results show that an Eos of *A. sativum*, *E. globulus*, and *L. angustifolia*, and the two chemicals deet and 2-undecanone were most effective against *L. decemlineata*. We also found that the most effective essential oils still showed elegant repellent effect when mingled to small concentrations. These essential oils can be suggested as adequate biopesticides agents under different controlled environmental conditions.

## Acknowledgments

This research was financially supported by the Operational Programme Integrated Infrastructure within the project: Sustainable smart farming systems taking into account the future challenges 313011W112, co-financed by the European Regional Development Fund.

## References

1. Abdalla MI, Abdelbagi AO, Hamma AMA, Laing MD. Use of volatile oils of garlic to control the cowpea weevil *Callosobruchus maculatus* (Bruchidae: Coleoptera). Scientific African Journal of Plant Soil. 2017;34:185. <https://doi.org/10.1080/02571862.2016.1225232>.
2. Abdel-Tawab H, Mossa SA. Green Pesticides: Essential Oils as Biopesticides in Insect-pest Management. Journal of Environmental Science and Technology. 2016;9:354-378. DOI: 10.3923/jest.2016.354.378.
3. Boukhatem MN, Boumaiza A, Nada HG, Rajabi M, Mousa SA. *Eucalyptus globulus* Essential Oil as a Natural Food Preservative: Antioxidant, Antibacterial and Antifungal Properties *in Vitro* and in a Real Food Matrix (Orangina Fruit Juice). Apply Science. 2020;10:5581. [htStps://doi.org/10.3390/app10165581](https://doi.org/10.3390/app10165581).
4. Chaudhari AK, Singh VK, Kedia A. Essential oils and their bioactive compounds as eco-friendly novel green pesticides for management of storage insect pests: prospects and retrospects. Environmental Science Pollution Resource. 2021;28:18918-18940. <https://doi.org/10.1007/s11356-021-12841-w>.
5. Chiam WY, Huang Y, Chen SX, Ho SH. Toxic and antifeedant effect of allyl disulphide on *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Sitophilus zeamais* (Coleoptera: Curculionidae). Journal of Economic Entomology. 1999;92(1):239-245. DOI: <https://doi.org/10.1093/jee/92.1.239>.
6. Conboy NJ, McDaniel T, George D. Volatile Organic Compounds as Insect Repellents and Plant Elicitors: an Integrated Pest Management (IPM) Strategy for Glass house Whitefly (*Trialeurodes vaporariorum*). Journal of Chemical Ecology. 2020;46:1090-1104. <https://doi.org/10.1007/s10886-020-01229-8>.
7. De Oliveira JL, Campos VR, Camara MC, Vechia JFD, De Matos STS, De Andrade DJ, *et al.* Hydrogels containing botanicals repellents encapsulated in zein nanoparticles for crop protection. American Chemical Society Appl. Nano Mater. 2019;8:33. <https://doi.org/10.1093/nar/gks1065>.
8. Dougoud J, Toepfer S, Bateman M, Jenner WH. Efficacy of homemade botanical insecticides based on traditional knowledge. A review. In Agronomy for Sustainable Development. 2019;39:1-22. DOI <https://doi.org/10.1007/s13593-019-0583>.
9. FAO. The State of Food and Agriculture, Overcoming water challenges in agriculture; c2020. p. 1564-3352 <https://doi.org/10.4060/cb1447en>.
10. Fusková M, Cagán E. Utilization of repellents for the control of Western corn rootworm (*Diabrotica virgifera* LeConte, 1868), Journal of Central European Agriculture. 2021;22(2):443-449. Received: January 21, 2021; accepted: March 29, 2021, DOI: /10.5513/JCEA01/22.2.3160.

11. Germinara GS, Di Stefano MG, De Acutis L, Pati S, Delfine S, De Cristofaro A, *et al.* Bioactivities of *Lavandula angustifolia* essential oil against the stored grain pest *Sitophilus granaries*. *Bull. Insectology*. 2017;70:129-139.
12. González-Coloma A, Reina M, Díaz CE, Fraga BM. Natural product based biopesticides for insect control. Mander HW, Lu (Eds.), *Comprehensive Natural Products II Chemistry and Biology*, Elsevier Ltd.; c2010. p. 237-268. <https://doi.org/10.1016/B978-008045382-8.00074-5>.
13. Grumezescu AM. New Pesticides and Soil sensors. *Nanotechnology in Agri - Food Industry*; c2017. ISBN 978 -0-12-804299-1.
14. Hin A. Colorado potato beetle management on potatoes: Current challenges and future prospects. *Fruit Vegetable Cereal Science and Biotechnology*. 2009;3(1):10-19. <https://doi.org/10.1080/15569543.2018.1554588>.
15. Isman MB. Botanical insecticides, deterrents and repellents in modern agriculture and an increasingly regulated world. *Annual Review Entomology*. 2006;51:45-66. Doi: 10.1146/annurev.ento.51.110104.151146.
16. Isman MB, Tak JH. Commercialization of insecticides based on plant essential oils: Past, Present, and Future. In *Green pesticides handbook: Essential oils for pest control*, Bioscience. Environment & Agriculture Boca Raton. 2017;7:27-39. DOI <https://doi.org/10.1201/9781315153131>.
17. Isman MB. Plant essential oils for pest and diseases management. In *Crop Protection*. 2000;19:603-8. DOI:[doi.org/10.1016/S0261-2194\(00\)00079-X](https://doi.org/10.1016/S0261-2194(00)00079-X).
18. Isman MB, Machal CM. Pesticides based on plant essential oils: from traditional practice to commercialization. Rai, M. C. Carpinella (Eds.), *Naturally Occurring Bioactive Compounds*, Elsevier; c2006. p. 29-44. [https://doi.org/10.1016/S1572-557X\(06\)03002-9](https://doi.org/10.1016/S1572-557X(06)03002-9).
19. Jahromi GM, Pourmirza AA, Safaralizadeh MH. Repellent effect of sirinol (garlic emulsion) against *Lasioderma serricorne* (Coleoptera: Anobiidae) and *Tribolium castaneum* (Coleoptera: Tenebrionidae) by three laboratory methods. *African Journal of Biotechnology*. 2012;11(2):280-288. DOI: <https://doi.org/10.5897/AJB11.2548>.
20. James C. Global Status of Commercialized Biotech/GM Crops; ISAAA: Ithaca, NY, USA, *Acquisition of Agri-biotech Applications* (ISAAA). 2011;20(50):1-135.
21. Kebede Y, Gebre-Michael T, Balkew M. Laboratory and field evaluation of neem (*Azadirachta indica* A. Juss) and chinaberry (*Melia azedarach* L.) oils as repellents against *Phlebotomus orientalis* and *P. bergeroti* (Diptera: Psychodidae) in Ethiopia. *Acta Tropica*. 2010;113:145-150. DOI: 10.1016/j.actatropica.2009.10.009.
22. Khallaayoune K, Biron JM, Chaoui A, Duvallat G. Efficacy of 1% geraniol (Fulltec<sup>®</sup>) as a tick repellent. *Parasite*. 2009;26:223-226. [http://dx.doi.org/10.1051/parasite/2009163223\\_2](http://dx.doi.org/10.1051/parasite/2009163223_2)
23. KOKO. Waraporn Juntarajumngong and Angsumarn Chandrapatya. Repellency, Fumigant and Contact Toxicities of *Litsea cubeba* (Lour.) persoon against *Sitophilus zeamais* Motschulsky and *Tribolium castaneum* (Herbst). *Kasetsart Journal Natural Sciences (Thailand)*. 2009;43(1):56-63.
24. Maia MF, Moore S. Plant - based insect repellents: a review of their efficacy, development and testing. In *Malaria Journal*. 2011;10(1):11-15. DOI 10.1186/1475-2875-10-S1-S11.
25. Moncada A, Obed C, Cruz N, Isabel X. Efecto, del-Tiempo, y. La dosis del *Allium sativum* L. en el Control Biologico del *Acanthoscelides Obtectus*. *Ingeniero Tesis. Facultad de Ingeniería y Arquitectura, Escuela Profesional de Ingeniería Ambiental; Universidad César Vallejo: Trujillo, Perú*; c2021. p. 60. Available online: [https://repositorio.ucv.edu.pe/bitstream/handle/20.500.12692/81158/Asencio\\_MCO\\_Navarro\\_CXI-SD.pdf?sequence=1&isAllowed=y](https://repositorio.ucv.edu.pe/bitstream/handle/20.500.12692/81158/Asencio_MCO_Navarro_CXI-SD.pdf?sequence=1&isAllowed=y) (accessed on 5 April 2021).
26. Mukarram M, Khan MMA, Zehra A, Choudhary S, Naeem M, Aftab T. Biosynthesis of lemongrass essential oil and the underlying mechanism for its insecticidal activity." In *Medicinal and Aromatic Plants*; c2021a. p. 429-443. Springer, Cham. [https://doi.org/10.1007/978-3-030-58975-2\\_18](https://doi.org/10.1007/978-3-030-58975-2_18).
27. Oerke E. Crop losses to pests. *The Journal of Agricultural Science*. 2006;144(1):31-43. Doi: 10.1017/S0021859605005708.
28. Panji HR. Some observations on the insecticidal activities of the fruit of Darek *Melia azedarach* (L). *Res. Bull. Punj. Univ. N. S. Sci*. 1964;15:4345-4346. <https://doi.org/10.1515/znc-2011-3-406>.
29. Plata-Rueda A, Martínez LC, Dos Santos M, Fernandes FL, Wilcken CF, Soares MA, Serrão JE, *et al.* Insecticidal activity of garlic essential oil and their constituents against the melanom beetle, *Tenebrio molitor* Linnaeus (Coleoptera: Tenebrionidae). *Scientific Reports*. 2017;7(1):464-466. DOI: <https://doi.org/10.1038/srep46406>.
30. Regnault R, Arnason V. Essential oils in insect control low - risk products in a high – stakes world. In *Annual Review of Entomology*. 2012;57:405-424. DOI: 10.1146/annurev-ento-120710-100554. Doi: 10.1146/annurev-ento-120710-100554.
31. Regnault-Roger C. Recherche de nouveaux biopesticides d'origine vegetalea caracte`reinsecticide: de: de: de: de: de marches me thodologiques et application aux plantes aromatiques me diterrane`ennes. *Biopesticides d'origine végétale* (2e éd.). 2008;98:25-50.
32. Regnault-Roger C. Essential oils in insect control KG. Ramawat and JM, Mérillon (Eds.). *Natural Products*, Springer-Verlag Berlin Heidelberg; c2013. p. 4087-4107. DOI: 10.1007/978-3-642-22144-6\_181.
33. Sanghong R, Junkum A, Chaithong U, Jitpakdi A, Riyong D, Tuetun B. Remarkable repellency of *Ligusticum sinense* (Umbelliferae), an herbal alternative against laboratory populations of *Anopheles minimus* and *Aedes aegypti* (Diptera: Culicidae). *Malaria Journal*. 2015;14(1):305-307. DOI: <https://doi.org/10.1186/s12936-015-0816-y>.
34. Shahriari M, Zibae A, Shamakhi L, Sahebzadeh N, Naseri D, Hoda H. Bio-efficacy and physiological effects of *Eucalyptus globulus* and *Allium sativum* essential oils against *Ephestia kuehniella* Zeller

- (Lepidoptera: Pyralidae). *Toxin Review*. 2019;39:422-433.
35. Tu H, Qin Y. Repellent effect of different celery varieties in *Bemisia tabaci* (Hemiptera: Aleyrodidae) Biotype, Q. *Journal of Economic Entomology*. 2017;110:1307-1316. DOI: 10.1093/jee/tox110.
36. Witting-Bissinger BE, Stumpf CF, Donohue KV, Apperson CS, Roe EM. Novel arthropod repellent BioUD, is an efficacious alternative to DEET. *Journal of Medical Entomology*. 2008;45(5):891-898. DOI: <https://doi.org/10.1603/0022-2585>.